

## Aspects of reproductive biology of the Greek fir (*Abies cephalonica* L.) in the Mt. Aenos National Park (Greece)

Patrizia Iossifina Politi<sup>1</sup>, Margarita Arianoutsou<sup>1</sup> and Kyriacos Georghiou<sup>2</sup>

<sup>1</sup> Department of Ecology and Systematics, Faculty of Biology, University of Athens

<sup>2</sup> Department of Botany, Faculty of Biology, University of Athens

Contact should be made to P.I. Politi, email [ppolith@biol.uoa.gr](mailto:ppolith@biol.uoa.gr)

### Keywords

*Abies cephalonica*; mediterranean forests; masting; seeds

### Aim/s and Introduction

*Abies cephalonica* L. is a forest tree species endemic to the mountainous regions of central and southern Greece. The National Park of Mount Aenos at the island of Cephalonia (western Greece), was founded in 1962 to ensure the protection and conservation of this unique *Abies cephalonica* insular population, covering an area of 28.620km<sup>2</sup>. This area runs almost 11km from NW to SE (38°09'04" N, 20°38'38" E) on a calcareous basement and reaches the height of 1628m above sea level. The current study is part of a Ph.D Thesis concerning the evaluation of the conservation status of the Greek fir in the Aenos National Park and it reports on aspects of the species' reproductive biology. The main objectives are to determine the extent of interannual variation in cone and viable seed production and to investigate a possible reproductive strategy employed by the species.

Previous research studies performed in the broader area of the Park have focused on botanical issues (e.g. Phitos and Damboldt 1985) and managing propositions, while the species has been studied from a genetic (Fady 1990 and 1991, Liu 1971) and silviculture point of view (Panetsos 1975). These works have indicated that although the species shows high levels of resistance to drought (comparing to other species of the genus *Abies* in the Mediterranean basin) its' current distribution on Mount Aenos has been severely reduced (Forestry Department of Kefalonia, 1996).

### Materials and Methods

Twenty sampling permanent plots, of 100m<sup>2</sup> each, were established at 11 different sites of the Park chosen in such a way so as to represent as much as possible of the landscape heterogeneity. The *Abies* population extends from 700 to 1620m height, although a few scattered individuals occur at lower elevations.

All mature trees occurring in the plots were mapped and measured. Every tree was monitored for its morphometric characteristics as well as for its cone and seed production on a three-year period.

Cones were manually collected (with a use of a crane) at maturity stage from the top of the trees and separately put into hard-paper boxes. The cones were measured and then left to dry in well ventilated rooms, where the exfoliation process was concluded and scales and seeds were separated from the central axis.

All cone morphometric characteristics (length, diameter, cone-axis) were measured in the lab, while cones, seeds and scales weights were subsequently weighted using an analytical balance (180g ± 0.1mg).

### Results

Flower buds usually open from late May to mid-June. Female flowers are born mostly in the top half of the crown and on the outer ends of branches. Male flowers are born in the lower half of the crown. Seed production appears to begin when trees have reached the age of 25-30 years old (> 5m long, > 9cm diameter at breast height).

Cone and viable seed production varied significantly between years. However, the timing of dispersal remained almost the same regardless of the level of seed production. Cones are formed on the trees late in spring, ripen in summer while seed dispersal takes place from middle October to early November.

The cones are erect, almost cylindrical, ending to a short-pointed apex of a yellowish green colour turning to brown at maturity. Their length is approximately 6.3±0.4cm, with a mean diameter of 4.6 ±0.1cm, while their mean weight varies between 65.3±3.4 g in years of low cone productivity and 74± 3.7g in the year of high cone productivity, as it is expressed by the number of cone bearing trees. The scales of the cones as well as the seeds on the same cone vary in size and weight, being smaller at the two edges and much bigger at the middle part of the cone where the sound seeds are located.

Measurements performed in 3 consecutive years revealed a high interannual variability in the cone production. Trees bearing cones in 2004 were only 4 out of 182 (0.0020/m<sup>2</sup>) in all permanent plots monitored, they reached the number of 5 (0.0025/m<sup>2</sup>) at the following year, while they were by far numerous in 2006, 99 individuals (0.049/m<sup>2</sup>) (Fig.1a). The average number of cones per tree seems to vary significantly too, being 16 ±3.4 in the year 2004, 19.4 ±3.4 in 2005 and 51.6 ±5.7 in 2006 (Fig. 1b).

As for the seed production, measurements performed in intact cones of the year 2005 indicated that each cone contains approximately 400 seeds. The number of sound seeds produced seems to vary also between years. Counting of seeds over a representative seedlot of 300 seeds selected from the central part of cones produced in 2005 from 30 individuals showed a 2% sound seeds and 3% of insect damaged seeds, while for 2006 a much higher percentage (40%) of sound seeds and a decrease of insect damage (0.3%) was revealed. The average weight of sound seeds was the same (0.8± 0.009g).

The average number of sound seeds offered as seed rain was 0.05/ m<sup>2</sup> at the year 2005 and increased to 98.80/ m<sup>2</sup> in the year of 2006.

## Discussion

During the period of the study a different reproductive pattern was recorded for the three consecutive years, possibly indicating a masting behaviour of *Abies cephalonica*. Mast-fruiting or masting behaviour has been interpreted mainly as the synchronous intermittent production of large seed crops in perennial plants (Kelly and Sork 2002) as well as the cumulative result of the reproductive patterns of individuals within a population (Koenig et al. 2003) in which there is a synchronization both in time and in size of crop.

The annual variability of cone production in populations of *Abies cephalonica* has been previously reported from observations from other parts of Greece (Panetsos 1975, Dafis 1986), but little is known about the production of seeds and sound seeds of the species.

Our data showed a significant concordance between yield of cone production and that of sound seed production. Similar trends have been reported by Houle (1999) for *Abies balsamea* in an old growth temperate forest of north-eastern North America.

Several hypotheses have been proposed with regard to the selective advantages of masting (Janzen 1971, Waller 1979, Smith et al. 1990, Lalonde and Roitberg 1992). Norton and Kelly (1988) and Kelly (1994) categorized these hypotheses into two types: (1) "resource matching," in which plants vary their reproductive effort in response to fluctuations in available resources; and (2) "economy of scale," in which larger reproductive effort is more efficient, favouring an occasional large effort rather than a regular, smaller one. However, there are still few studies with sufficient quantitative data to test these hypotheses for masting species.

Our study needs further observations and analysis in order to fully justify this pattern as well to classify *Abies cephalonica* in one of the above categories.

## References

- Dafis S. 1986. Forest Ecology. Ghiahoudi and Ghiapouli publications. Thessaloniki. Greece.
- Fady B. 1990. Genetic variability of height growth components of the Greek fir (*Abies cephalonica*). Canadian Journal of Forest Research 20:1453-1460.
- Fady B. 1991. Variability of juvenile greek firs and stability of characteristics with age. Silva Genetica 40:91-99.
- Forestry Department of Kefalonia. 1996. General management plan of the National Park of Mount Aenos. Kefalonia, Greece.
- Houle G. 1999. Mast seeding in *Abies balsamea*, *Acer saccharum* and *Betula alleghaniensis* in an oldgrowth, cold temperate forest of north-eastern North America. Journal of Ecology 87:413-422.
- Janzen, D. 1971. Seed predation by animals. Annual Review of Ecology and Systematics 2:465-492.
- Kelly D. 1994. The evolutionary ecology of mast seeding. Trends in Ecology & Evolution 9:465-470.
- Kelly D and Sork V. 2002. Mast seeding in perennial plants: Why, How, Where? Annual Review of Ecology and Systematics 33:427-447.
- Koenig W, Kelly D, Sork V, Duncan R, Elkinton J, Peltonen M and Westfall R. 2003. Dissecting components of population-level variation in seed production and the evolution of masting behavior. Oikos 102:581-591.
- Lalonde R and Roitberg B. 1992. On the evolution of masting behavior in trees: predation or weather? American Naturalist 139:1293-1304.
- Liu T. 1971. A monograph of the genus *Abies*. National Taiwan University, College of Agriculture, Department of Forestry, Taipei, Taiwan, China.
- Norton D and Kelly D. 1988. Mast seeding over 33 years by *Dacrydium cupressinum* Lamb. (rimu) (Podocarpaceae) in New Zealand: the importance of economies of scale. Functional Ecology 2:399-408.
- Panetsos K. 1975. Monograph of *Abies cephalonica* Loudon. Yugoslavian Academy of Science, Annales Forestales 11:1-22.
- Phitos D and Damboldt J. 1985. Die flora der insel of Kefallinia. Botanika Chronica. University of Patras. Patras. Greece.
- Smith C, Hamrick J and Kramer C. 1990. The advantage of mast years for wind pollination. American Naturalist 136:154-166.
- Waller D. 1979. Models of mast fruiting in trees. Journal of Theoretical Biology 80:223-232.




Figure 1 – a. Percentage of mature trees bearing cones per sampling plot (100m<sup>2</sup>) and b. average number of cones per tree, during the years of study.