Ecology of Southern African Sandveld Lizards (Lacertidae, *Nucras*) Martin H. van der Meer¹, Martin J. Whiting^{1,2}, and William R. Branch³

Sandveld lizards (genus Nucras) are widespread in southern Africa, but are generally secretive and poorly known. We examined 385 preserved specimens from five species of Nucras collected over a broad time span (104 years) and a geographic area covering most of South Africa and Swaziland. We had three main objectives: to test for sexual size dimorphism, to quantify male and female reproductive cycles, and to determine diet. In addition, we examined the importance of scorpions in the diet of Nucras based on previous studies reporting an unusually high incidence of scorpions in the diet of Kalahari N. tessellata. Males of all species except N. lalandii had significantly larger heads than females of the same body size, although females had significantly greater snout-vent length than males in three of the five species examined. The general reproductive pattern was for females to contain vitellogenic eggs during late spring and early summer. No females simultaneously contained oviductal and vitellogenic eggs, which suggests that females produce only one clutch per season. Clutch size was unrelated to female body size in all species examined. Female follicle volume generally coincided with male testicular volume, indicating for the most part synchronized reproductive cycles between the sexes. Testis volume was generally highest during spring-early summer, with only N. holubi showing a second peak in autumn. We recorded 15 arthropod orders in the diet of Nucras. All species feed on invertebrates, primarily insects, and, to a lesser degree, spiders and centipedes. Termites, grasshoppers, and beetles both numerically and volumetrically dominated their diet. We found no evidence that scorpions form a major part of the diet of any of these five species of Nucras, including N. tessellata from biomes outside the Kalahari Desert.

NDERSTANDING broad-scale ecological patterns of diverse living organisms is fundamental to developing life history theory (Stearns, 1976). In this regard, ecological studies of lizards have been extremely fruitful (Milstead, 1967; Huey et al., 1983; Pianka, 1986; Vitt and Pianka, 2005). For example, extensive ecological data sets combined with phylogenies of major lizard clades have facilitated our understanding of the role of early evolutionary events on the evolution of ecological traits and the associated present day structuring of lizard communities (Vitt et al., 2003; Vitt and Pianka, 2005). However, even the combined lifetime efforts of two key lizard ecologists, E. R. Pianka and L. J. Vitt, have resulted in ecological data for only about 4.25% of 'lizards' and 2.2% of squamates (Vitt and Pianka, 2007). Furthermore, these data are biased by region. We need additional ecological data from more taxa, and from taxa occupying a greater variety of habitat types in which different predator assemblages and resources occur (Vitt and Pianka, 2007) in order to elucidate and understand broad-scale ecological patterns.

One of the best-studied ecological attributes of lizards is their diet. Previous studies have focused on lizard foraging mode, preference for particular prey, and how they detect and capture prey (Vitt and Pianka, 2005; Cooper, 2007; Reilly et al., 2007). Many lizard species are generalized insectivores that consume prey of a range of sizes, while others are highly specialized. For example, *Ameiva ameiva* will eat many small prey and some large ones (reviewed in Vitt and Pianka, 2007), while *Moloch horridus* has a highly specialized diet consisting of ants only (Pianka and Pianka, 1970). Because most lizards thus far studied are generalists and eat a wide variety of prey sizes, any exceptions to this pattern are of particular interest to ecologists. Of specific interest are lizards that eat large prey and occupy a 'top predator' position in a food web.

The southern African lacertid Nucras tessellata is a 'top predator' (Huey et al., 2001) that consumes a high proportion of scorpions (53%), considered to be large and energetically rich prey (Pianka, 1986). Nucras tessellata is also an active forager that retrieves scorpions from their day-time retreats (Pianka, 1986). Current knowledge of the ecology of Nucras (Sandveld lizards) in general is restricted to N. tessellata and N. intertexta from the Kalahari Desert (Huey and Pianka, 1977; Pianka et al., 1979; Huey and Pianka, 1981; reviewed in Pianka, 1986). Using museum specimens, we studied the ecology of five species of the lizard genus Nucras with three main objectives: to test for sexual size dimorphism, to quantify male and female reproductive ecology, and to examine diet. With respect to diet, our primary aim was to determine the extent to which these species of Nucras eat scorpions, including N. tessellata from areas outside of the Kalahari Desert.

MATERIALS AND METHODS

Study system.—Lacertids of the genus Nucras are mainly restricted to scrubland or savannah habitats in southern Africa, with isolated species occurring in Angola (*N. scalaris*) and East Africa (*N. boulengeri*). The genus was last revised by Broadley (1972), who recognized seven species, some containing additional races. A molecular phylogeny of *Nucras* is currently in preparation (Bauer, Branch, and Burger, unpubl.), with indications that additional cryptic taxa may be subsumed within *N. tessellata* (Branch, 2006). We have assigned specimens to species as accepted in Branch (1998), but are aware that Kalahari arenicolous populations of *N. tessellata* and *N. intertexta* have been the subject of ecological study; both are active foragers and *N.*

¹School of Animal, Plant and Environmental Sciences, University of the Witwatersrand, Private Bag 3, Wits 2050, South Africa; E-mail: (MVDM) martinhvandermeer@gmail.com.

² Department of Brain, Behaviour and Evolution, Macquarie University, Sydney, NSW 2109, Australia; E-mail: martin.whiting@mq.edu.au. Send reprint requests to this address.

³ Port Elizabeth Museum, P.O. Box 13147, Humewood 6013, South Africa, and Research Associate, Department of Zoology, Nelson Mandela Metropolitan University, Port Elizabeth, South Africa; E-mail: wrbranch@bayworld.co.za.

Submitted: 20 July 2009. Accepted: 13 July 2010. Associate Editor: J. D. Litzgus.

^{© 2010} by the American Society of Ichthyologists and Herpetologists 🖨 DOI: 10.1643/CE-09-134

tessellata specializes largely on scorpions (reviewed in Pianka, 1986; also see Huey and Pianka, 1977; Huey et al., 1984). We present baseline ecological and morphological data from museum specimens for five of the ten currently recognized species of *Nucras* from eastern and southern Africa.

Sexual size dimorphism.—We examined 385 specimens, including N. holubi (77), N. intertexta (76), N. lalandii (82), N. ornata (84), and N. tessellata (66). The following morphological variables were measured to test for sexual size dimorphism: head length, width, and height to the nearest 0.01 mm using digital calipers; snout-vent length (SVL) and tail (original, complete) length, both measured to the nearest 1 mm. All data met assumptions of normality and homoscedacity. Sexual size dimorphism was tested using a one-factor ANOVA between mean values for adult males vs. females. We also tested for heterogeneity of slopes. If slopes did not differ significantly an ANCOVA was performed. Finally, we used a reduced major axis (RMA) regression (Fairbairn, 1997; Bohonak and van der Linde, 2004) to investigate whether growth patterns of characters differed significantly between the sexes.

Reproduction.—We made a mid-ventral incision to expose the gonads in order to score sex and reproductive condition. Females were considered sexually mature if the oviducts were muscular and scored as sexually active at the time of preservation if eggs or large ovarian follicles were present. We recorded the diameter of the largest vitellogenic follicle or oviductal egg to the nearest 0.01 mm using digital calipers. Males were considered sexually mature if testes were enlarged/turgid (immature males had flat, ribbon-like testes), and defined as sexually active at time of preservation if the epididymides were convoluted. Testes were measured for length and width to the nearest 0.01 mm using digital calipers. We calculated egg and testis volume using the formula for a prolate spheroid (Vitt et al., 1993):

Volume =
$$4/3\pi$$
 (length/2) * (width/2)².

Diet.—We determined diet by removing and examining stomach contents of preserved specimens. Prey items were spread out on a petri dish and identified to order. We also measured the length and width of each intact prey item to the nearest 0.01 mm using digital calipers, and calculated prey volume using the formula for a prolate spheroid (see above).

Niche breadth measure for numerical and volumetric diet data were estimated using the reciprocal of Simpson's (1949) formula:

$$\mathbf{B}=1/\sum_{i=1}^n p_i^2,$$

where i = resource category, n = total number of categories, and p = proportion of resource category i. Values range from 1 (exclusive use of one prey type) to n (even use of all prey types). Prey categories are represented by the arthropod orders of prey items found in stomachs.

We also calculated percent utilization for each prey type numerically and volumetrically (the pi \times 100 of Pianka, 1973, 1986). A regression analysis was used to determine the relationship between SVL and mean prey volume, and SVL and number of prey items consumed. We also compared

prey size among species by performing an ANOVA on the number of prey items and prey volume/individual. A Bonferroni *post-hoc* test was used to determine sources of differences. Analyses were performed with SAS Enterprise Guide 3.0 (SAS Institute Inc., 2007), and all means are reported \pm 1 SE. We used parametric tests when the assumptions of normality and homoscedasticity could reasonably be met.

RESULTS

Sexual size dimorphism.—Summary statistics of morphological measurements for the five species are given in Table 1, and frequency of SVL in Figure 1. Overall, sexual dimorphism in SVL and head size was variable among species and in both directions (males or females larger). Females of Nucras holubi and N. lalandii were significantly larger in SVL than males; likewise, N. intertexta females had significantly larger SVL, head length, and width than males. Nucras ornata males had significantly wider and deeper heads than females (Table 2). When the effect of body size was removed, N. holubi males had significantly larger heads than females (length, width, and height), N. intertexta males had significantly longer and deeper heads than females, N. lalandii males had significantly longer heads than females, N. ornata males had significantly wider and deeper heads than females, and N. tessellata males had significantly longer heads than females (Table 2). For most species, the head typically grew isometrically in relation to SVL (Table 3).

Tail length differed significantly among species ($F_{4,231} = 34.203, P < 0.001$). A Bonferroni *post-hoc* test showed that *N*. *lalandii*, *N*. *ornata*, and *N*. *tessellata* had proportionately longer tails followed by *N*. *intertexta* and *N*. *holubi*.

Reproduction.—The smallest male and female showing evidence of sexual maturity were 59 and 61 mm (N. holubi), 67 and 65 mm (N. intertexta), 69 and 71 mm (N. lalandii), 57 and 59 mm (N. ornata), and 63 and 61 mm (N. tessellata), respectively (Fig. 1). Females collected during the breeding season contained clutches varying from 1–7 eggs (N. holubi: n = 12, mean = 4.08 ± 0.5), 4–8 eggs (N. intertexta: n = 12, mean = 6 ± 0.41), 4–14 eggs (*N. lalandii*: n = 2, mean = $7 \pm$ 1), 4–13 eggs (*N. ornata*: n = 4, mean = 5.5 \pm 0.87), and 3–7 eggs (N. tessellata: n = 6, mean = 4.67 \pm 0.56). Clutch size was not related to female body size (SVL), although this is likely due to small sample sizes: N. holubi ($r^2 = 0.039$, P =0.54), N. intertexta ($r^2 = 0.011$, P = 0.75), N. ornata ($r^2 =$ 0.041, P = 0.70), N. lalandii ($r^2 = 0.01723$, P = 0.35), and N. tessellata ($r^2 = 0.061$, P = 0.59). Egg size increased significantly with SVL for N. holubi ($r^2 = 0.189$, P < 0.05) and *N. tessellata* ($r^2 = 0.323$, P < 0.01), but not for *N*. intertexta ($r^2 = 0.051$, P = 0.24), N. lalandii ($r^2 = 0.006$, P =0.68), or *N. ornata* ($r^2 = 0.031$, P = 0.38). Clutch size did not differ significantly among species when controlling for body size $(F_{4,43} = 2.49, P = 0.06)$, although this marginally nonsignificant difference could be an artifact of sample size. Seasonal variation in ovarian volume of the largest follicle indicated that females were gravid during spring-summer and autumn (N. holubi), summer (N. intertexta [one individual in autumn], N. lalandii, N. tessellata), and spring-summer (N. ornata, Fig. 2). No evidence of double clutching was found in specimens examined.

Body size (SVL) was significantly correlated with testis volume for *N. holubi* ($r_p = 0.54$, P < 0.001, n = 51), *N.*

b B L	
lle	
tai	
°.	
Å	ŝ
p	ter
JC -	ne
.⊑.	1
/eľ	E
s	.⊑
tail	are
fe	ţ
ple	len
E	en
č	sur
Ę.	ea
s	E
ard	Π
Ľ.	Ċ
È	30
ō	Ш
es.	٦, f
Jes	-
nt	ll E
are	<u> </u>
ď	ata
.⊑.	elle
ar	ess
ges	l. te
ang	<
1	32
S	II
+1	, f
fe	16
oc	II
Je l	<u> </u>
Ire	ťa
IS 8	'na
ear	0
ž	2
ď.	7
s sl	ī
ra	÷
μ	27,
Ę.	Ш
ls o	3
lua	1
-i	ŭ
ġ	a
ē	>
Ę	<u>_</u>
Ma	28
≧	Ш
ua	Ъ,
ě	-
£	
Š	드
isti	xta
feri	rte
raci	nte
haı	۲. i
Ū	< '
<u>ica</u>	36
log	Ш
q	, f
Jrp	
ĕ	l
	느
e 1	iq
ab	olt
	<u> </u>

			Morphological traits		
Species/sex	SVL	Tail length	Head length	Head width	Head height
N. holubi					
Females $(n = 19)$ Males $(n = 45)$	$63.95 \pm 1.65 (50-77)$ $59.98 \pm 0.80 (49-72)$	$117.07 \pm 4.94 (87-166)$ $112.49 \pm 2.35 (83-134)$	$12.75 \pm 0.36 (9.76-15.38)$ $13.05 \pm 0.21 (9.78-15.71)$	8.74 ± 0.25 (7.15-10.77) 8.94 ± 0.14 (6.85-10.51)	$\begin{array}{l} 6.76 \pm 0.22 \; (5.43 - 8.72) \\ 6.81 \pm 0.12 \; (5.11 - 8.41) \end{array}$
N. intertexta					
Females $(n = 25)$ Males $(n = 38)$	73.32 ± 1.77 (56–91) 63.32 ± 0.98 (49–77)	147.47 ± 5.46 (112–186) 135.62 ± 3.59 (97–176)	$13.23 \pm 0.28 (11.08 - 15.82)$ $12.54 \pm 0.17 (10.28 - 15.09)$	$6.47 \pm 0.19 (3.38-8.04)$ $6.30 \pm 0.14 (8.57-11.76)$	$9.04 \pm 0.21 (6.88 - 10.91)$ $8.44 \pm 0.13 (6.95 - 10.74)$
N. lalandii					
Females $(n = 36)$	$89.11 \pm 1.62 (71-107)$	$159.78 \pm 5.20 (104-207)$	$14.21 \pm 0.22 (11.26 - 16.38)$	10.49 ± 0.19 (8.01-13.54)	$7.80 \pm 0.14 (6.07 - 9.17)$
N. ornata	(101-14) 64.7 - 16.00	(EUZ-EE) 10.0 - 00.001	(07:01-+6:11) 07:0 - 00:01	(a/.11-10.0) 17.0 - an.n1	(10.6-10.9) 01.0 - 70.1
Females $(n = 25)$ Males $(n = 48)$	$74.52 \pm 2.55 (47-103)$ $77.75 \pm 2.11 (47-111)$	$147.50 \pm 5.68 (107-207)$ 159.24 + 5.71 (104-231)	$13.36 \pm 0.43 (8.25 - 16.76)$ 14 51 + 0 38 (9 98 - 20 83)	$10.06 \pm 0.33 (7.15 - 13.49)$	7.45 ± 0.26 (4.69–9.94) 8 40 + 0.24 (5.29–12.24)
N. tessellata					
Females $(n = 18)$ Males $(n = 37)$	$71.00 \pm 1.84 (58-86)$ 69.08 $\pm 1.07 (52-83)$	162.73 ± 6.11 (114–196) 165.42 ± 5.42 (98–219)	$13.01 \pm 0.26 (11.04 - 14.30)$ $13.25 \pm 0.19 (10.04 - 15.79)$	9.04 ± 0.23 (7.59-10.88) 8.72 ± 0.15 (6.01-10.39)	$\begin{array}{l} 6.26 \pm 0.24 \ (4.31 - 8.08) \\ 6.08 \pm 0.11 \ (4.21 - 7.54) \end{array}$

intertexta ($r_p = 0.51$, P < 0.001, n = 46), N. lalandii ($r_p = 0.54$, P < 0.001, n = 37), N. ornata ($r_p = 0.65$, P < 0.001, n = 44), and weakly for N. tessellata ($r_p = 0.30$, P = 0.05, n = 42). Testes were most turgid during spring–early summer, with a second peak in autumn (N. holubi), spring–summer (N. intertexta), summer (N. lalandii, N. ornata [one individual in autumn]), and in N. tessellata did not show a clear pattern (Fig. 3). Male and female reproductive cycles appeared to be generally synchronous.

Diet.—Of the 385 lizard stomachs examined, 167 (43%) were empty (26/77 = 34%, N. holubi; 28/76 = 37%, N. intertexta; 37/82 = 45%, N. lalandii; 44/84 = 52%, N. ornata; 32/66 = 48%, N. tessellata). The remaining 218 stomachs contained 1229 prey items. A wide variety of prey taxa were consumed, including termites, spiders, scorpions, insect eggs and in a few cases, lizard tails (Table 4). For all five species the largest percentage of individual prey items consumed were termites. However, the largest volume of stomach contents varied by species and constituted termites (39%, N. holubi), grasshoppers (15%, N. intertexta; 63%, N. lalandii; 50%, N. ornata), and ants (53%, N. tessellata). Mean number of prey items per stomach and average volume of prey per stomach were 10.61 ± 2.42 and 444.59 ± 79.46 mm³ (*N. holubi*), 5.98 \pm 2.03 and 529.66 \pm 79.46 mm³ (*N. intertexta*), 5.82 \pm 1.78 and 579.40 \pm 107.66 mm³ (N. lalandii), 4.53 \pm 1.66 and $631.52 \pm 172.73 \text{ mm}^3$ (N. ornata), and 4.51 ± 1.33 and $830.86 \pm 273.60 \text{ mm}^3$ (*N. tessellata*), respectively.

No significant relationship was found between SVL and number of prey (*N. holubi*: $r_s = 0.035$, P = 0.19; *N. intertexta*: $r_s = 0.03$, P = 0.24; *N. lalandii*: $r_s = 0.007$, P = 0.58; *N. ornata*: $r_s = 0.02$, P = 0.35; *N. tessellata*: $r_s = 0.002$, P = 0.79). Mean prey volume was significantly correlated with SVL for *N. intertexta* ($r_s = 0.484$, P < 0.001, n = 48) and *N. ornata* ($r_s = 0.196$, P < 0.005, n = 40), but not for *N. holubi* ($r_s = 0.005$, P = 0.64, n = 51), *N. lalandii* ($r_s = 0.06$, P = 0.10, n = 45) and *N. tessellata* ($r_s = 0.005$, P = 0.69, n = 34).

Niche breadths for prey number ranged from 1.29 (*N. holubi*)–3.40 (*N. lalandii*), indicating that many individuals of just a few prey types (termites, grasshoppers, and ants) were consumed (Table 4). For *N. holubi* and *N. intertexta*, niche breadths for prey volume were high enough to suggest that several prey categories were important, while in *N. lalandii*, *N. ornata*, and *N. tessellata* niche breadths for prey volume were low and suggest that only a few prey categories were important (Table 4).

When controlling for body size, no significant differences were found between sexes for total prey number ($F_{1,217} = 0.14$, P = 0.71), total prey volume ($F_{1,217} = 2.19$, P = 0.14), and mean prey volume/individual ($F_{1,217} = 0.24$, P = 0.63). Likewise, when body size was controlled for, no difference was found among species for total prey number ($F_{4,217} = 1.69$, P = 0.15) and total prey volume ($F_{4,217} = 0.93$, P = 0.45). However, mean prey volume/individual was significantly ($F_{4,217} = 8.74$, P < 0.001) different among species. A Bonferroni *post-hoc* test showed that *N. ornata* consumed the largest prey items, followed by *N. lalandii* and *N. tessellata*, *N. holubi*, and *N. intertexta* (Fig. 4).

DISCUSSION

Of the five species examined, females were the larger (SVL) sex in three species and males in two species, suggesting that different selective pressures are at play, depending on

th (V.



Fig. 1. Snout–vent length (mm) frequency for male and female *Nucras* spp. Solid arrows indicate minimum female size at sexual maturity; open arrows indicate minimum male size at sexual maturity. A = N. *holubi*, B = N. *intertexta*, C = N. *lalandii*, D = N. *ornata*, E = N. *tessellata*.

species. Unfortunately we did not separately measure trunk size, which appears to be under stronger selection relative to overall body size in certain female lacertids from Europe (Braña, 1996). In general, female lizards have relatively longer trunks than males as a consequence of fecundity selection favoring individuals with a capacity to carry more eggs or offspring, while males frequently have larger heads because of contest competition (Cooper and Vitt, 1989; Braña, 1996; Olsson et al., 2002), although other factors also contribute to SSD (Cox et al., 2003). When the effect of body size was removed, some aspect of head size in males of all species except N. lalandii was significantly larger than that of females. Although head size is typically acted on by sexual selection for contest competition (Olsson et al., 2002; Husak et al., 2009), intrasexual niche divergence can explain differences in head size as a result of sex-specific trophic ecology (Shine, 1989; Herrell et al., 1999). While there are some differences in types of prey eaten by males and females of different Nucras spp., prey number, total prey volume, and prey volume/individual did not differ significantly between sexes. Therefore, the most parsimonious mechanism explaining head size sexual dimorphism is male contest competition, and not trophic partitioning, but other factors could also be important (Cox et al., 2003). Sexual dimorphism in southern African lacertids is largely unstudied, although female Pedioplanis burchelli have larger bodies than males, but smaller heads (Nkosi et al., 2004). Among eight species of European lacertids (Podarcis, Lacerta) males have larger heads than females, but females have larger abdomens, although overall body size could be significantly greater in either sex (Braña, 1996). Abdomen length increases at a faster rate relative to SVL in females, and this is explained by fecundity selection. Conversely, larger male head size was most likely due to sexual selection with no support for intersexual dietary niche divergence. In general, lacertids appear to conform to a typical pattern seen in lizards: fecundity selection acting on female abdomen size and sexual selection acting on head size (Braña, 1996).

Females contained vitellogenic eggs during late spring and early summer. No females simultaneously contained oviductal and vitellogenic eggs, which suggests that females of all species produce only one clutch per season, although N. holubi and N. intertexta show some evidence of autumn breeding. Clutch size for all species combined ranged from 1–14 eggs, with means ranging between 4–6 (excluding N. *lalandii*). Pianka (1986) reports a mean clutch size of 3.3 \pm 0.7 SD for eight *N. tessellata*, while we report a mean clutch size of 4.7 for six individuals. Clutch size and maternal body size were not significantly related, although small sample sizes for most taxa likely obscured any real relationships. Overall, male Nucras typically had enlarged testes during spring and summer, followed by regression in autumn and winter. In female Nucras, follicles were typically enlarged during spring-summer, followed by regression in late summer and continuing through winter (but see N. holubi, Fig. 2). Therefore, male and female reproductive cycles are largely synchronous.

Stomachs we examined from museum specimens were collected over a broad time span (104 years) and a geographic area covering most of South Africa and Swaziland. Therefore, the samples should be representative of dietary preference. *Nucras* were largely insectivorous, but also included spiders and centipedes in their diet. Their diet was numerically dominated by termites and volumetrically

Table 2. Sexual Size Dimorphism in Morphological Traits for *Nucras* spp. Results of a one-factor ANOVA between mean values for adult males vs. females, the tests for heterogeneity of slopes for *N. holubi* (n = 64), *N. intertexta* (n = 63), *N. lalandii* (n = 57), *N. ornata* (n = 73), and *N. tessellata* (n = 55), followed by ANCOVA if slopes did not differ significantly. Only lizards with complete tails were included for tail length (see Table 1). Sex was used as the factor in all analyses. Asterisk refers to a significant (P < 0.05) difference between mean values for adult males vs. females. The M/F column indicates which of the sexes has the larger trait.

			ANOVA	L Contraction of the second seco		Heterog	geneity o	of slopes test		Intercep	t	
Species	Trait	F	df	Р	M/F	F	df	Р	F	df	Р	M/F
N. holubi	SVL	5.83	1,63	0.02*	F							
N. intertexta		28.56	1,62	0.0001*	F							
N. lalandii		9.16	1,56	0.004*	F							
N. ornata		0.62	1,72	0.43								
N. tessellata		0.92	1,54	0.34								
N. holubi	Tail length	0.90	1,51	0.35		0.68	1,50	0.41	0.04	1,50	0.84	
N. intertexta		3.48	1,43	0.07		0.01	1,42	0.92	0.006	1,42	0.94	
N. lalandii		0.02	1,44	0.88		0.04	1,43	0.85	3.19	1,43	0.08	
N. ornata		1.64	1,48	0.21		0.09	1,47	0.77	1.41	1,47	0.24	
N. tessellata		0.09	1,45	0.76		0.92	1,44	0.34	1.15	1,44	0.29	
N. holubi	Head length	0.55	1,63	0.46		0.32	1,63	0.57	31.35	1,63	0.0001*	Μ
N. intertexta		5.15	1,62	0.03*	F	0.22	1,62	0.64	9.78	1,62	0.03*	Μ
N. lalandii		2.40	1,56	0.13		5.76	1,56	0.02	—	—	_	
N. ornata		3.60	1,72	0.06		1.24	1,72	0.27	3.11	1,72	0.08	
N. tessellata		0.54	1,54	0.47		0.52	1,54	0.47	4.78	1,54	0.03*	Μ
N. holubi	Head width	0.61	1,63	0.44		0.74	1,63	0.39	25.52	1,63	0.001*	Μ
N. intertexta		6.69	1,62	0.01*	F	0.17	1,62	0.68	6.30	1,62	0.01*	Μ
N. lalandii		2.21	1,56	0.14		1.15	1,56	0.29	0.02	1,56	0.89	
N. ornata		4.65	1,72	0.03*	Μ	0.12	1,72	0.73	9.20	1,72	0.003*	Μ
N. tessellata		1.45	1,54	0.23		0.38	1,54	0.54	0.58	1,54	0.45	
N. holubi	Head height	0.06	1,63	0.82		2.53	1,63	0.12	8.02	1,63	0.006*	Μ
N. intertexta		0.57	1,62	0.45		0.01	1,62	0.91	3.69	1,62	0.06	
N. lalandii		1.39	1,56	0.24		0.75	1,56	0.39	0.03	1,56	0.86	
N. ornata		6.03	1,73	0.02*	Μ	0.07	1,72	0.79	16.57	1,72	0.0001*	Μ
N. tessellata		0.64	1,54	0.43		0.88	1,54	0.35	0.02	1,54	0.88	

 Table 3.
 Results of a Reduced Major Axis (RMA) Regression Showing the Relationship between the Growth Rate of the Head (Height, Length, Width) in Relation to SVL.

		Males			Females	;
	r^2	Slope	95% Cl	r ²	Slope	95% Cl
N. holubi						
Head height Head length Head width	0.8064 0.8846 0.8969	1.164 1.089 1.051	1.018–1.309 0.9836–1.194 0.955–1.147	0.5571 0.8371 0.7154	1.316 1.155 1.101	0.9381-1.694 0.954-1.356 0.8478-1.355
N. intertexta						
Head height Head length Head width	0.2744 0.7133 0.7017	1.382 0.9097 0.9854	1.029–1.736 0.7635–1.056 0.8217–1.199	0.29 0.7907 0.8139	1.196 0.7903 0.9152	0.7982–1.594 0.6476–0.9331 0.7593–1.071
N. lalandii						
Head height Head length Head width	0.4221 0.5674 0.5095	0.8252 0.8139 0.8884	0.6272-1.023 0.645-0.9829 0.692-1.085	0.546 0.7408 0.5624	0.986 0.8799 0.9961	0.7643–1.208 0.7304–1.029 0.7762–1.216
N. ornata						
Head height Head length Head width	0.8806 0.5765 0.7931	1.08 0.8885 1.1	0.9755–1.185 0.7258–1.051 0.9589–1.24	0.8781 0.3729 0.8907	1.07 0.9061 0.9414	0.9283–1.212 0.6336–1.179 0.8232–1.06
<i>N. tessellata</i> Head height Head length Head width	0.6182 0.7715 0.5932	1.028 0.8502 1.004	0.8249–1.231 0.7203–0.98 0.7998–1.209	0.6399 0.7938 0.7929	1.28 0.9203 1.055	0.9401–1.619 0.7355–1.105 0.8429–1.268



Fig. 2. Volume of largest follicle for Nucras spp. in relation to time of the year. Numbers adjacent to points indicate the number of overlapping values. A = N. holubi, B = N. intertexta, C = N. lalandii, D = N. ornata, E = N. tessellata.

Fig. 3. Volume of largest testis for Nucras spp. in relation to time of the year. Numbers adjacent to points indicate the number of overlapping values. A = N. holubi, B = N. intertexta, C = N. lalandii, D = N. ornata, E = N. tessellata.

:

Dec

Dec

:

		N. hc	lubi (n =	51)			N. inter	texta (n =	48)			N. Ialo	ndii (n = .	45)			N. orna	ta (n = 4	(0			N. tess	ellata (n :	= 34)	
		%		%			%		%			%		%			%		%			%		%	
Prey type	No.	No.	Vol.	Vol.	Freq.	No.	No.	Vol.	Vol.	Freq.	No.	No.	Vol.	Vol.	Freq. 1	٩o.	No.	Vol.	Vol.	Freq.	No.	No.	Vol.	Vol.	Freq.
Orthoptera	20	3.8	5372.7	24.6	16	14	4.9	3916.3	15.4	13	57	21.8	16496.4	63.3	36.0	28	15.7 1	2549.9	49.9	25	19	12.1	3862.6	13.7	18
Blattaria	9	1.1	541.4	2.5	ß	9	2.1	3304.5	13.0	ъ						2	1.1	120.0	0.5	2					
Mantodea	-	0.2	274.5	1.3	-	-	0.3	79.7	0.3	-															
Isoptera	468	87.8	8596.7	39.3	17	195	67.9	1931.7	7.6	œ	94	36.0	860.9	3.3	4.0 1	16	65.2	2141.1	8.5	4	95	60.5	3196.6	11.3	9
Coleoptera	12	2.3	3420.6	15.6	Ø	20	7.0	2911.3	11.5	15	13	5.0	4733.1	18.2	5.0	-	0.6	275.3	1.1	2	б	5.7	4623.7	16.4	Ŋ
Hemiptera	2	0.4	179.4	0.8	-	10	3.5	1116.6	4.4	м						-	0.6	103.1	0.4	-	2	1.3	158.7	0.6	2
Lepidoptera	М	0.6	546.6	2.5	Ю	6	3.1	2981.5	11.7	ſ						М	1.7	335.9	1.3	2	2	1.3	459.7	1.6	2
Neuroptera																					ß	3.2	493.1	1.8	4
Diptera	ы	0.6	726.4	3.3	7	Ю	1.0	236.4	0.9	ю						Ю	1.7	678.0	2.7	Ю					
Hymenoptera	2	0.4	225.8	1.0	2	2	0.7	138.7	0.5	2						-	0.6	197.6	0.8	-					
Ants		0.2	49.9	0.2	-				0.0							-	0.6	73.3	0.3	-	18	11.5	4921.6	53.0	4
Insect eggs											88	33.7	376.1	1.4	4.0										
Araneae	10	1.9	420.0	1.9	Ø	19	6.6	1710.0	6.7	14	ß	1.9	374.8	1.4	5.0	13	7.3	3855.4	15.3	10	9	3.8	360.2	1.3	ß
Solifugae	-	0.2	273.4	1.2	-	-	0.3	1253.8	4.9	-															
Scorpiones	2	0.4	359.3	1.6	2	-	0.3	698.2	2.7	-	2	0.8	2740.2	10.5	2.0	2	1.1	360.1	1.4	2					
Chilopoda		0.2	763.9	3.5	-	ß	1.7	2562.4	10.1	ß		0.4	335.0	1.3	1.0	9	3.4	4422.8	17.6	4		0.6	89.4	0.3	
Diplopoda											-	0.4	133.4	0.5	1.0	-	0.6	49.9	0.2	-					
Lizard tail	-	0.2	128.3	0.6	-	-	0.3	2582.7	10.2	-															
Total	533	100	21879	100	69	287	100	25423.8	100	77	261	100	26049.8	100	58 1	78 1	00 2	5162.4 1	00	58	157 1	00	8165.6 1	00	47
Niche																									
breadths		1.29		4.10			2.10		9.66			3.40		2.24			2.19		3.20			2.50		2.94	

Table 4. Summary of Diet for *Nucras* spp. Insect taxonomy (prey categories) follows Borror et al. (1992). Frequency is the number of lizards containing a specific prey category in the respective sample. All Lepidoptera were larval. No. is the total number of individual prey items from a particular arthropod group, followed by its percentage (% No.) as a proportion of all identified arthropod prey. Vol. is the number of all interviews of a narricular arthropod group, while & volume of all interviews of a narricular arthropod group in the proportion for the number of hole of total number of individual previews of a narricular arthropod group, followed as a percentage (% No.) as a proportion of all identified arthropod prev. Vol. is the number of all interviews of a narricular arthropod group in the proportion for total volume of all interviews of a narricular arthropod group will be proportion for total volume of a narrow items of a narricular arthropod group will be proportion to total volume of a narrow items of a narrow items of a narrow items of a narrow items of a narrow item of a narrow items of a narrow items of a narrow items of a narrow item of a narrow item of a narrow item of a narrow items of a narrow item of a narrow item of a narrow items of a narrow item of a na



Fig. 4. Distribution of individual prey sizes (volume) consumed by *Nucras* spp. A = N. *holubi*, B = N. *intertexta*, C = N. *lalandii*, D = N. *ornata*, E = N. *tessellata*.

dominated by termites, grasshoppers, and beetles. All five species had similarly high proportions of termites and orthopterans in their diet, suggesting that although they eat a wide variety of prey, they are quick to take advantage of high quality prey items. Typically, widely foraging lacertids eat relatively slow-moving prey such as termites or inactive prey such as scorpions (Huey and Pianka, 1981). Termites are an important dietary component for many South African lizards (Pianka, 1986; Bauer et al., 1990; Shuttleworth et al., 2008), including other Nucras (Branch, 1998) and desert dwelling lizards (Huey and Pianka, 1981). However, termites are patchily distributed both spatially and temporally in arid areas of southern Africa (Lepage and Darlington, 2000; Traniello and Leuthold, 2000). Because all Nucras are likely to be active foragers (Cooper and Whiting, 1999), they may gain substantial nourishment from excavating subterranean termites or by encountering termite foraging parties or swarms. Termites, when present, were in high numbers (up to 70) in the gut, suggesting that lizards binge-feed on termites when they are available.

More than half the lizards examined (57%) were in a positive energy balance (i.e., stomachs contained food). In a global review of the proportion of lizards containing gut contents, Huey et al. (2001) found that of 101 diurnal lizard species, only $10.5 \pm 9.6\%$ had empty stomachs, and when only wide-foraging lizards were included, this number increased slightly to $15.1 \pm 1.3\%$ (n = 53 spp.). Therefore, the Nucras we examined had a lower frequency of individuals in positive energy balance, although the possibility that collectors kept individuals alive for sufficient time for their stomach contents to be digested, can not be discounted. Interestingly, Pianka (1986) reports that the scorpion specialist N. tesselata from the Kalahari Desert had a frequency of 16.1% of individuals with empty stomachs. In our limited sample of N. tesselata (n = 34), only one individual (2.9%) had an empty stomach.

Nucras, like other lacertids, are diurnal foragers (Pianka et al., 1979; Pianka, 1986). Many prev items found would need to be retrieved from beneath surface objects or sand because they are not diurnally surface active (hidden prey). Such prey include antlions, centipedes, scorpions, and even some spiders. Lizards would likely have to use vomerolfaction rather than visual searching, to find them, and Cooper (1990) has suggested that teiids and lacertids use chemical cues to identify and locate non-moving or hidden prey. However, in the case of *N. tesselata* from the Kalahari, it is possible that they locate scorpion burrows visually before entering them to retrieve the scorpion (R. Huey, pers. comm.). All of the species in our study consumed spiders and centipedes; however, N. ornata appeared to rely quite heavily on these as one of its main food sources (33% by volume). Millipedes (Diplopoda) were found in the gut of a single N. lalandii and a single N. ornata. Millipedes are known to produce toxins from repugnatory glands (Vitt, 1992) and are thus avoided by most lizards (Vitt and Cooper, 1986; Wapstra and Swain, 1996). Active foragers can detect released toxins and avoid them (see Evans, 1961); however, as only two lizards consumed one millipede each, they may have done so accidentally or these millipedes might be less toxic.

Nucras tessellata from the Kalahari Desert represent a paradox among lacertids because they consume relatively low numbers of termites (4.6% by volume) but high numbers of scorpions (53% by volume; Pianka et al., 1979;

Huey and Pianka, 1981) in a termite-rich landscape where other lacertids eat large quantities of termites (Pianka, 1986). In contrast, we found no scorpions in the 34 N. tessellata we examined from outside the Kalahari, and they contained a relatively higher volume of termites (11.4%). There are several possible explanations for this finding. First, N. tessellata from the Kalahari may be specifically distinct from other populations of currently recognized N. tessellata (Bauer, Branch, and Burger, unpubl.); therefore, dietary differences may be an artifact of phylogeny. Second, N. tessellata from outside the Kalahari occur in very different habitats that are sometimes less sandy and more rocky. This habitat may make scorpion capture more difficult. Third, N. tessellata from the Kalahari are active at high temperatures and begin activity later in the morning than other lacertids (Huey and Pianka, 1977; Pianka et al., 1979), under conditions that may preclude termite activity or when termites are further underground. Therefore, they may simply not eat termites because the prevailing thermal conditions make termites less accessible. While these theories are somewhat speculative, they are testable. We need field studies of Nucras to assess the relative influence of phylogeny and environment on ecology and life history. In particular, understanding how Kalahari N. tessellata have become such dietary specialists compared to closely related congeners will contribute to our understanding of the evolution of feeding specialization in lizards.

MATERIAL EXAMINED

TMP specimen numbers: 94, 96, 562, 563, 567, 683, 1144, 1385, 1389, 1390, 1505, 1764, 2217, 2501, 2537, 3169, 3170, 3171, 3426, 3675, 3723, 4431, 4433, 4434, 4435, 4502, 4522, 4526, 4539, 9220, 11055, 11068, 11133, 11181, 11187, 11188, 11334, 11628, 12289, 12290, 12352, 12376, 12812, 15676, 15705, 15707, 15951, 18030, 18056, 18179, 19377, 20129, 21130, 22180, 27161, 27166, 27840, 27841, 28791, 28820, 29997, 30184, 30434, 33292, 33420, 33539, 33540, 33541, 33632, 34024, 34052, 34140, 35292, 35404, 35497, 35693, 35734, 36045, 36133, 36282, 36393, 36621, 36694, 37211, 37212, 37213, 37214, 37218, 37219, 37220, 37716, 37717, 37718, 37844, 37845, 37846, 37847, 37898, 39102, 39500, 39920, 42459, 42460, 42461, 43259, 43285, 43288, 43289, 43291, 43292, 43293, 43304, 43306, 43307, 43308, 43308, 43309, 43310, 43311, 43393, 43407, 43411a, 43412, 43428, 43446, 43450, 43474, 43476, 43515, 43518, 43559, 43560, 43561, 43562, 43576, 43669, 43671, 43769, 47458, 47580, 49504, 50111, 50325, 51562, 51563, 51564, 51565, 51568, 52450, 52451, 53294, 53909, 53910, 53922, 54768, 55223, 55301, 55554, 55714, 55752, 55761, 55773, 55909, 55913, 56041, 56100, 57525, 62994, 63006, 63986, 66485, 66499, 66766, 66865, 66970, 67345, 67945, 68838, 68839, 68840, 68848, 69068, 70049, 70631, 70942, 70943, 70947, 71071, 72340, 72342, 72343, 72344, 72348, 72350, 72351, 72352, 72353, 72354, 72356, 72359, 72658, 72659, 72661, 72663, 72666, 72667, 72669, 72673, 72674, 72675, 72676, 72677, 72678, 72679, 72684, 72686, 72687, 72689, 72690, 72691, 72693, 72694, 72695, 72696, 72698, 72699, 72708, 76868, 76869, 76870, 76871, 76872, 76873, 76874, 76877, 76878, 76879, 76880, 76881, 76882, 76883, 76884, 76885, 77797, 77802, 77811, 77812, 77828, 77830, 77834, 77836, 77837, 77838, 77839, 77840, 77841, 77842, 77843, 77844, 77845, 77846, 77849, 77850, 77851, 77853, 77854, 77857, 77858, 77859, 77860, 77862, 77863, 77865, 77867, 77867, 77868, 77869, 77870, 77871, 77873, 77875, 77876, 77877, 77879, 77880, 77881, 77882, 77886, 77887, 77888, 77889, 77890, 77891, 77892, 77894, 77900, 78316, 78317, 78318, 78322, 78324, 78325, 78326, 78327, 78330, 78331, 78332, 78338, 78669, 78670, 78671, 78705, 78706, 78708, 79455, 80027, 80036, 80070, 83011, 83016, 83370, 83563, 83564, 83566, 83662, 83831, 84030, 84729, 84861, 84983, 85378, 85426. PEM specimen numbers: 569, 1914, 1939, 2378, 2693, 2985, 3043, 3044, 3053, 4099, 4101, 4132, 4135, 4142, 4147, 4220, 4294, 4857, 4980, 5027, 5067, 5079, 5791, 5906, 7064, 7070, 7102, 7155, 7156, 7207, 7247, 7278, 7386, 7590, 7629, 7681, 7899, 8055, 8056, 8147, 8164, 8168, 8400, 8421, 8438, 8439, 8440, 8450, 8478, 8483, 8719, 9707, 10425, 10426, 10427, 10428, 10429, 10430, 10431, 10432, 10433, 10438, 10439, 10440, 10441, 10442, 10454, 10455, 10456, 10457, 10458, 10459, 10460, 10461, 10462, 10463, 10464, 10465, 10466, 10467, 10468, 10469, 10470, 10471, 10472, 10473, 10474, 10475, 10476, 10477, 10478, 10479, 10480, 10481, 10482, 10483, 10484, 10679, 10680, 11111, 12161, 12162, 12410, 13259, 13344, 13355, 13357, 13358, 13373, 13507, 13762, 15990, 15991, 15992, 15993, 15994, 15995, 15996, 15997, 15998, 15999, 16000, 16001, 16002, 16003, 16004, 16005, 16006, 16007, 16008, 16009, 16010, 16011, 16012, 16013, 16014, 16015, 16016, 16017, 16018, 16019, 16020, 16021, 16022, 16023, 16024, 16025, 16026, 16027, 16028, 16029, 16030, 16031, 16032, 16033, 16034, 16035, 16036, 16037, 16038, 16039, 16040, 16041, 16042, 16043, 16044, 16492, 16493, 16495, 16496, 16497, 16498, 16499, 16500, 16503, 16504, 16506, 16872, 16873, 16874.

ACKNOWLEDGMENTS

We thank L. Mashini and L. Mahlangu (Northern Flagship Institution, formerly Transvaal Museum) for the loan of specimens and H. van der Meer for many hours of data capture. R. Huey and E. Pianka are thanked for providing useful comments that greatly improved the quality of this manuscript.

LITERATURE CITED

- Bauer, A. M., A. P. Russell, and B. D. Edgar. 1990. Utilization of the termite *Hodotermes mossambicus* (Hagen) by gekkonid lizards near Keetmanshoop, South West Africa. South African Journal of Zoology 24:239–243.
- Bohonak, A. J., and K. van der Linde. 2004. RMA: Software for Reduced Major Axis regression for Java. http://www. kimvdlinde.com/professional/rma.html
- Borror, D. J., C. A. Triplehorn, and N. F. Johnson. 1992. An Introduction to the Study of Insects. Saunders College Publishers, New York.
- **Braña**, F. 1996. Sexual dimorphism in lacertid lizards: male head increase vs female abdomen increase? Oikos 75: 511–523.
- **Branch, W. R.** 1998. Field Guide to the Snakes and Other Reptiles of Southern Africa. Third edition. Struik Publishers, Cape Town.
- Branch, W. R. 2006. Priorities for systematic studies on southern African reptiles, p. 2–20. *In*: A Plan for Phylogenetic Studies of Southern African Reptiles. W. R. Branch, K. A. Tolley, M. Cunningham, A. M. Bauer, G. Alexander, J. A. Harrison, A. A. Turner, and M. F. Bates (eds.). Proceedings of a workshop held at Kirstenbosch, February

2006. Biodiversity Series No. 5. South African National Biodiversity Institute, Pretoria.

- **Broadley**, D. G. 1972. A review of the *Nucras tessellata* group (Sauria: Lacertidae). Arnoldia Rhodesia 5:1–36.
- **Cooper, W. E., Jr.** 1990. Prey odor detection by teiid and lacertid lizards and the relationship of prey odor detection to foraging mode in lizard families. Copeia 1990:237–242.
- Cooper, W. E., Jr. 2007. Lizard chemical senses, chemosensory behavior, and foraging mode, p. 237–270. *In*: Lizard Ecology: The Evolutionary Consequences of Foraging Mode. S. M. Reilly, L. D. McBrayer, and D. B. Miles (eds.). Cambridge University Press, Cambridge.
- **Cooper, W. E., Jr., and L. J. Vitt.** 1989. Sexual dimorphism of head and body size in an iguanid lizard: paradoxical results. The American Naturalist 133:729–735.
- Cooper, W. E., Jr., and M. J. Whiting. 1999. Foraging modes in lacertid lizards from southern Africa. Amphibia– Reptilia 20:299–311.
- Cox, R. M., S. L. Skelly, and H. B. John-Alder. 2003. A comparative test of adaptive hypotheses for sexual size dimorphism in lizards. Evolution 57:1653–1669.
- **Evans**, L. T. 1961. Structure as related to behavior in the organizations of populations of reptiles, p. 148–178. *In*: Vertebrate Speciation. W. F. Blair (ed.). University of Texas Press, Houston.
- Fairbairn, D. J. 1997. Allometry for sexual size dimorphism: patterns and processes in the coevolution of body size in males and females. Annual Review of Ecology and Systematics 28:659–687.
- Herrel, A., L. Spithoven, R. V. Damme, and F. D. Vree. 1999. Sexual dimorphism of head size in *Gallotia galloti*: testing the niche divergence hypothesis by functional analyses. Functional Ecology 13:289–297.
- Huey, R. B., A. F. Bennett, H. John-Alder, and K. A. Nagy. 1984. Locomotor capacity and foraging behaviour of Kalahari lacertid lizards. Animal Behaviour 32:41–50.
- Huey, R. B., and E. R. Pianka. 1977. Seasonal variation in thermoregulatory behavior and body temperature of diurnal Kalahari lizards. Ecology 58:1066–1075.
- Huey, R. B., and E. R. Pianka. 1981. Ecological consequences of foraging mode. Ecology 62:991–999.
- Huey, R. B., E. R. Pianka, and T. W. Schoener. 1983. Lizard Ecology: Studies of a Model Organism. Harvard University Press, Cambridge.
- Huey, R. B., E. R. Pianka, and L. J. Vitt. 2001. How often do lizards "run on empty"? Ecology 82:1–7.
- Husak, J. F., A. Kristopher Lappin, S. F. Fox, and J. A. Lemos-Espinal. 2009. Bite-force performance predicts dominance in male venerable collared lizards (*Crotaphytus antiquus*). Copeia 2006:301–306.
- Lepage, M., and J. P. E. C. Darlington. 2000. Population dynamics of termites, p. 333–361. *In*: Termites: Evolution, Sociality, Symbioses, Ecology. T. Abe, D. E. Bignell, and M. Higashi (eds.). Kluwer Academic Publishers, Dordrecht, Netherlands.
- Milstead, W. W. 1967. Lizard Ecology: A Symposium. University of Missouri Press, Columbia, Missouri.
- Nkosi, W. T., N. J. L. Heideman, and J. H. Van Wyk. 2004. Reproduction and sexual size dimorphism in the lacertid lizard *Pedioplanis burchelli* (Sauria: Lacertidae) in South Africa. Journal of Herpetology 38:473–480.

- Olsson, M., R. Shine, E. Wapstra, B. Ujvari, and T. Madsen. 2002. Sexual dimorphism in lizard body shape: the roles of sexual selection and fecundity selection. Evolution 56:1538–1542.
- Pianka, E. R. 1973. The structure of lizard communities. Annual Review of Ecology and Systematics 4:53–74.
- Pianka, E. R. 1986. Ecology and Natural History of Desert Lizards. Princeton University Press, Princeton, New Jersey.
- Pianka, E. R., and H. D. Pianka. 1970. The ecology of Moloch horridus (Lacertilia: Agamidae) in the Western Australian Desert. Emu 70:17–22.
- Pianka, E. R., R. B. Huey, and L. R. Lawlor. 1979. Niche segregation in desert lizards, p. 67–115. *In*: Analysis of Ecological Systems. D. J. Horn, G. R. Stairs, and R. D. Mitchell (eds.). Ohio State University Press, Columbus, Ohio.
- **Reilly, S. M., L. B. McBrayer, and D. B. Miles (eds.).** 2007. Lizard Ecology: The Evolutionary Consequences of Foraging Mode. Cambridge University Press, Cambridge.
- Shine, R. 1989. Ecological causes for the evolution of sexual dimorphism: a review of the evidence. Quarterly Review of Biology 64:419–461.
- Shuttleworth, C., P. le Fras, N. Mouton, and J. H. van Wyk. 2008. Group size and termite consumption in the armadillo lizard, *Cordylus cataphractus*. Amphibia–Reptilia 29:171–176.
- Simpson, E. H. 1949. Measurement of diversity. Nature 163:688.
- **Stearns, S.** C. 1976. Life history tactics: a review of the ideas. Quarterly Review of Biology 51:3–47.
- Traneillo, J. F. A., and R. H. Leuthold. 2000. Behavior and ecology of foraging in termites, p. 141–168. *In*: Termites: Evolution, Sociality, Symbioses, Ecology. T. Abe, D. E. Bignell, and M. Higashi (eds.). Kluwer Academic Publishers, Dordrecht, Netherlands.
- Vitt, L. J. 1992. Mimicry of millipedes and centipedes by elongate terrestrial vertebrates. Research and Exploration 8:76–95.
- Vitt, L. J., and W. E. Cooper, Jr. 1986. Foraging and diet of a diurnal predator (*Eumeces laticeps*) feeding on hidden prey. Journal of Herpetology 20:208–415.
- Vitt, L. J., and E. R. Pianka. 2005. Deep history impacts present day ecology and biodiversity. Proceedings of the National Academy of Sciences of the United States of America 102:7877–7881.
- Vitt, L. J., and E. R. Pianka. 2007. Feeding ecology in the natural world, p. 141–172. *In*: Lizard Ecology: The Evolutionary Consequences of Foraging Mode. S. M. Reilly, L. D. McBrayer, and D. B. Miles (eds.). Cambridge University Press, Cambridge.
- Vitt, L. J., E. R. Pianka, W. E. Cooper, Jr., and K. Schwenk. 2003. History and the global ecology of squamate reptiles. The American Naturalist 162:44–60.
- Vitt, L. J., P. A. Zani, J. P. Caldwell, and J. P. Durtsche. 1993. Ecology of the whiptail lizard *Cnemidophorus deppii* on a tropical beach. Canadian Journal of Zoology 71:2391–2400.
- Wapstra, E., and R. Swain. 1996. Feeding ecology of the Tasmanian skink, *Niveoscincus ocellatus* (Squamata: Scincidae). Australian Journal of Zoology 44:205–213.