

Somatostatin receptor SPECT

Giovanna Pepe, Roy Moncayo, Emilio Bombardieri & Arturo Chiti

**European Journal of Nuclear
Medicine and Molecular Imaging**


ISSN 1619-7070

Volume 39

Supplement 1

Eur J Nucl Med Mol Imaging (2012)
39:41-51

DOI 10.1007/s00259-011-2019-2



Your article is protected by copyright and all rights are held exclusively by Springer-Verlag. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your work, please use the accepted author's version for posting to your own website or your institution's repository. You may further deposit the accepted author's version on a funder's repository at a funder's request, provided it is not made publicly available until 12 months after publication.

Somatostatin receptor SPECT

Giovanna Pepe · Roy Moncayo · Emilio Bombardieri ·
Arturo Chiti

Published online: 3 March 2012
© Springer-Verlag 2011

Abstract Somatostatin is a peptide with a broad distribution in the nervous system and acts as a neurotransmitter in several organs, having a wide range of mainly inhibiting effects, such as the suppression of growth hormone release, as well as the inhibition of pancreatic and gastrointestinal hormone release. Five somatostatin receptor subtypes have been cloned and demonstrated to have an emphasized expression in all human tumours. In particular, type 2 receptors were identified as the most frequently represented on the surface of neuroendocrine tumour cells, providing the molecular basis for many clinical applications of somatostatin analogues. Towards the end of the 1980s, the *in vivo* demonstration of somatostatin receptors on the surface of some tumours raised interest in receptor imaging, and indeed the peptide receptor overexpression on tumour cells, as compared to normal tissues, constitutes the basis for molecular imaging of these tumours. This review intends to illustrate the development of single photon emission radiopharmaceuticals for the study of somatostatin receptors and their application in diagnostic imaging.

Keywords Somatostatin · Somatostatin receptor scintigraphy (SRS) · Single photon emission computed tomography (SPECT) · Neuroendocrine tumours (NET) · Gastroenteropancreatic (GEP) tumours · Peptide receptor radionuclide therapy (PRRT)

G. Pepe · A. Chiti (✉)
Nuclear Medicine, IRCCS Humanitas,
Milan, Italy
e-mail: arturo.chiti@cancercenter.humanitas.it

R. Moncayo
Nuclear Medicine, Medical University of Innsbruck,
Innsbruck, Austria

E. Bombardieri
Nuclear Medicine, National Cancer Institute,
Milan, Italy

Introduction

Somatostatin is a peptide with a broad distribution in the nervous system and acts as a neurotransmitter in several organs, having a wide range of mainly inhibiting effects, such as the suppression of growth hormone release, as well as the inhibition of pancreatic and gastrointestinal hormone release [1, 2]. Five somatostatin receptor subtypes have been cloned of whichsstr 1 and 4 are grouped into one family andsstr 2, 3 and 5 into another. They all are G protein-coupled receptors located at the cell membrane [3].

Pangerl et al. [4] reported that onlysstr3 seems to have an emphasized expression in all human tumours, whilstsstr2 appears to be the most frequently represented receptor over the surface of neuroendocrine tumour (NET) cells [5], providing the molecular basis for many clinical applications of somatostatin analogues [6].

Towards the end of the 1980s, the *in vivo* demonstration of somatostatin receptors on the surface of some tumours raised interest in receptor imaging [7], and indeed the peptide receptor overexpression on tumours cells, as compared to normal tissues [8, 9], constitutes the basis for molecular imaging of these tumours.

This review intends to be a “voyage” through the development of single photon emission radiopharmaceuticals for the study of somatostatin receptors and their application in diagnostic imaging.

We will describe the somatostatin analogues and the derived radiopharmaceuticals, describe the imaging technique and show the results and applications in the clinical scenario.

Somatostatin analogues

Somatostatin is an acid polypeptide broadly distributed throughout different organs and tissues including the central

nervous system. Structural and functional studies on this peptide demonstrated the amino acid components that are considered fundamental for the peptide activity in the body; therefore, a number of peptides showing similarities in the binding affinity of the native somatostatin have been synthesized [10].

The most important octapeptides in clinical use are octreotide and lanreotide, registered in several countries for clinical use [11] (Table 1).

Octreotide-based radiopharmaceuticals

The radiopharmaceutical that was first used in patients to study human NET is ^{123}I -Tyr³-octreotide, a radioiodinated somatostatin analogue with a Tyr substitution [12, 13]. Due to the costs of producing ^{123}I , the need of special technology and skills for peptide iodination and the difficult interpretation of images due to the accumulation of the radiopharmaceutical in the bowel—a consequence of the predominantly biliary clearance of the product—led to the development of ^{111}In -DTPA-D-Phe¹-octreotide (also called ^{111}In -pentetreotide). This tracer became the first radioreceptor imaging tracer commercially available (OctreoScan, Mallinckrodt Medical, St. Louis, MO, USA) that was approved in 1994 by the US Food and Drug Administration as an imaging agent for somatostatin receptor-positive NET. ^{111}In -pentetreotide has shown high accuracy for imaging NET [14–16] (Fig. 1).

Due to the affinity of ^{111}In -pentetreotide to bind to sstr2 it has shown effectiveness in diagnosing and localizing NET. By means of this diagnostic ability staging of the disease through the detection of metastases has been made possible [17, 18].

As compared to ^{123}I -Tyr-octreotide, it shows less intestinal accumulation as it is mainly cleared via the kidneys [19] and appears to be more suitable for late imaging thanks to its longer half-life, improving the interpretation of the scanning of the upper abdomen.

Given the importance and success of $^{99\text{m}}\text{Tc}$ as a routine isotope in nuclear medicine imaging, it came

as a logical consequence that $^{99\text{m}}\text{Tc}$ -labelled somatostatin analogues have been developed, namely the $^{99\text{m}}\text{Tc}$ -N- α -(6-hydrazinonicotinoyl)-octreotide ($^{99\text{m}}\text{Tc}$ -EDDA/HYNIC OCT) [20, 21] (Fig. 2). At that time propagation of the tracer was hampered by proprietary rights on the peptide. For this reason it has remained a tracer for in-house use. Interestingly it is currently being offered as a routine product by POLATOM (Warsaw, Poland). Besides imaging of NET it has been used in the study of thyroid-associated orbitopathy [22].

Further improvements in the development of radio-receptor radiopharmaceuticals have been achieved with the introduction of DOTA a more universal chelator for metal ions, in the attempt to develop peptides that could be labelled to other isotopes such as ^{90}Y with therapeutic rather than diagnostic aims. On the diagnostic side, ^{111}In -DOTA-D-Phe¹-Tyr³-octreotide shows similar bio-distribution of ^{111}In -DTPA-D-Phe¹-octreotide [23].

Three other tracers are ^{111}In -DOTANOC (1-Nal³-octreotide) [24], ^{111}In -DOTANOC-ATE (1-Nal³-Thr⁸-octreotide) and ^{111}In -DOTABOC-ATE, (Bz-Thi³-Thr⁸-octreotide), the applications of which are limited though and are not currently commercially available [25, 26]. The replacement of octreotide with octreotate has eventually shown a decrease of lipophilicity [27], boosting the affinity for the sstr2.

Among octreotate-based radiopharmaceuticals there are maltotriose- ^{123}I -Tyr³-octreotate (^{123}I -Mtr-TOCA) that however did not show major imaging advantages [28] and $^{99\text{m}}\text{Tc}$ -Demotate for which preclinical evaluations are available [29–31].

It is interesting to mention that this Demotate peptide has been further developed into a positron emission tomography (PET) tracer in a more sophisticated combination with receptor-based reporter gene features [32].

The most recent contribution to $^{99\text{m}}\text{Tc}$ -labelled peptides is Demotensin used for the detection of brain metastases [33].

Other analogue-based radiopharmaceuticals

The successful introduction of ^{111}In -pentetreotide as a radiopharmaceutical for scintigraphic imaging coupled with the knowledge about other somatostatin analogues with different affinity profiles led as a consequence to efforts in developing new radiopharmaceuticals with higher sensitivity or a wider somatostatin receptor subtype affinity profile.

Some of these somatostatin analogues are characterized by the macrocyclic chelator DOTA instead of DTPA in their structure, suitable for the labelling not only with ^{111}In for diagnostic use but also with beta emitters, namely ^{90}Y and ^{177}Lu , used for peptide receptor radionuclide therapy (PRRT).

An example is ^{111}In -DOTA-lanreotide, known also as MAURITIUS (Multicenter Analysis of a Universal

Table 1 sstr subtype selectivity to endogenous somatostatin and somatostatin analogues modified from Volante et al. [10]

Agonist	<i>K_i</i> (nM)				
	sstr1	sstr2	sstr3	sstr4	sstr5
SS-14	1.1	1.3	1.6	0.53	0.9
SS-28	2.2	4.1	6.1	1.1	0.07
Octreotide	>1,000	0.6	34.5	>1,000	7
Lanreotide	>1,000	0.8	107	>1,000	5.2
Vapreotide	>1,000	5.4	31	45	0.7
SOM-230	9.3	1.0	1.5	>100	0.2




Fig. 1 Imaging with ^{111}In -pentetreotide. Patient affected by well-differentiated pancreatic NET. Diagnostic scan pre-therapy showed advanced disease with nodal, liver and bone metastases. After two

cycles of treatment (one with ^{177}Lu -DOTA-TATE and one with ^{90}Y -DOTA-TATE) the post-therapy scan showed good response to treatment




Fig. 2 Imaging with $^{99\text{m}}\text{Tc}$ -EDDA/HYNIC-TOC. Case of unknown primary. SRS imaging detected focal uptake in the ileum and a mediastinal node. CT fusion and registration was performed between separate scans

Receptor Imaging and Treatment Initiative: a European study), that has been used in Europe for dosimetry studies for therapy with ^{90}Y -DOTALAN. Compared to ^{111}In -DTPA-D-Phe¹-octreotide, it was shown to have lower affinity for NET but high affinity for intestinal adenocarcinoma and differentiated thyroid cancer [34]. One drawback of ^{111}In -DOTA-lanreotide, however, is its affinity for bone marrow which makes diagnostic interpretation difficult.

$^{99\text{m}}\text{Tc}$ -Depreotide, P829, commercially also known as NeoTect or NeoSpect, was approved for lung cancer studies [35]. Its applications for NET in the abdomen are reduced by the short half-life of the labelling isotope, thus not permitting late imaging to reduce the abdominal background [36].

$^{99\text{m}}\text{Tc}$ -Vapreotide (RC-160 Octastatin) was developed for applications in gastroenterology and NET. It shows high binding affinity for sstr 2 and 5 and less for sstr 3 and 4 [37] (Table 2).

Somatostatin receptor scintigraphic imaging

Technique

Revised somatostatin receptor scintigraphy (SRS) guidelines have recently been published by the European Association of Nuclear Medicine (EANM) [38]. In an abridged way the method can be summarized as follows.

Previous to the moment of injecting the tracer preparation of the patient is needed. Any “cold” somatostatin analogues should be suspended 4 weeks before scintigraphy. In addition the administration of a mild oral laxative to reduce the abdominal interferences is recommended.

The preferred administered activity is 200 MBq [39–42] and a peptide amount of 10 µg, which is not expected to produce significant pharmacological effects.

The radiopharmaceutical is prepared according to the manufacturer’s indications and undergoes quality controls during which the activity is measured in a calibrated ionization chamber and the radiochemical purity is verified with a thin-layer chromatography (TLC) method.

A gamma camera equipped with a medium-energy parallel-hole collimator with ^{111}In photopeaks (172 and 245 keV) set for 20% windows is used to acquire 4-, 24- and also 48-h post-injection planar total body scans. Further single photon emission computed tomography (SPECT) images of the area of interest, neck, thorax or abdomen can also be acquired at both 4 and 24 h, but preferably at 24 h.

As we further explain in the following section, CT coregistration has been reported to improve the localization of the lesions, thanks to the attenuation correction [43–45].

The $^{99\text{m}}\text{Tc}$ -labelled octreotide has also been used for image fusion studies [46].

Images are processed according to the preferences of each nuclear medicine department. Image viewing depends on the technology available on site.

Normal findings and imaging pitfalls

Thyroid, spleen, liver, kidneys and sometimes pituitary gland are visualized as normal foci of uptake in SRS [19, 47]. The uptake in pituitary, thyroid, spleen and adrenals depends on receptor binding. The excretion via the liver may cause the visualization of bowel; therefore, the use of laxatives prior to late scanning (at 24–48 h) is recommended

Table 2 Somatostatin receptor radiopharmaceuticals

Radiopharmaceutical		Availability
Full name	Abbreviation	
^{123}I -Tyr ³ -octreotide		Historical
^{111}In -DTPA-D-Phe ¹ -octreotide	^{111}In -DTPA-OCT (^{111}In -pentetreotide)	CCA
^{111}In -DOTA-D-Phe ¹ -Tyr ³ -octreotide	^{111}In -DOTA-TOC	In house
^{111}In -DOTA-Tyr ³ -octreotate	^{111}In -DOTA-TATE	In house
^{111}In -DOTA-NaI ³ -octreotide	^{111}In -DOTA-NOC	In house
^{111}In -DOTA-NaI ³ -Thr ⁸ -octreotide	^{111}In -DOTANOC-ATE	Research
^{111}In -DOTA-Bz-Thi ³ -Thr ⁸ -octreotide	^{111}In -DOTABOC-ATE	Research
$^{99\text{m}}\text{Tc}$ -EDDA/HYNIC-Tyr ³ -octreotide	$^{99\text{m}}\text{Tc}$ -EDDA/HYNIC-TOC	In house
$^{99\text{m}}\text{Tc}$ -EDDA/HYNIC-Tyr ³ -octreotate	$^{99\text{m}}\text{Tc}$ -EDDA/HYNIC-TATE	In house
^{123}I -Mtr-TOCA-octreotate		Research
$^{99\text{m}}\text{Tc}$ -Demotate		In house
^{111}In -DOTA-lanreotide	^{111}In -DOTA-LAN (MAURITIUS)	In house
$^{99\text{m}}\text{Tc}$ -depreotide		CCA
$^{99\text{m}}\text{Tc}$ -vapreotide		Research

CCA currently commercially available

in order to differentiate aspecific areas of uptake [48]. The radiopharmaceutical is also excreted via the kidneys and is reabsorbed in the renal tubular cells: on the one hand this causes the possible visualization of the urinary bladder and on the other a higher radiation exposure of the kidneys, requiring a good hydration status and even infusion at therapeutically administered doses for renal protection.

A specific uptake in breast or in sites of recent surgery, the visualization of gallbladder or the presence of accessory spleens could lead to misinterpretation (Table 3).

There are also some other possible pitfalls emanating from other conditions such as inflammations and infections, granulomatous diseases, nodular goiter and cerebrovascular accidents [49].

Some of the SRS drawbacks related to the limited spatial resolution and the lack of anatomical landmarks may be overcome by the use of hybrid SPECT/low-energy CT integrated diagnostic systems. There is an interesting number of studies reporting an improvement in terms of diagnostic accuracy in general nuclear medicine when using a SPECT/CT device, and in particular the contribution of CT fusion to NET imaging has been reported by Krausz et al. [50] who found an incremental value of SPECT/CT over SPECT alone of 32%, with a subsequent change of patient management in 14% of cases. Similar results were described by Hillel et al. [44].

In a small series of patients studied by Moreira et al. [51] CT fusion imaging proved to be of additional value in NET imaging, and more recently Perri et al. [52] prospectively compared the performance of ^{111}In -pentetreotide SPECT and SPECT/CT in patients with known or suspected NET and analysed the results on a patient-by-patient and lesion-by-lesion basis. SPECT/CT showed a 95.3% sensitivity and 92.1% specificity whilst for SPECT alone sensitivity was 95.3% and specificity was 71%, confirming that CT fusion improving the anatomical localization of the areas of uptake is useful in reducing equivocal findings.

Table 3 Causes of potential misinterpretation of negative results with ^{111}In -pentetreotide scintigraphy modified from Kwekkeboom et al. [54]

Cause
Possible competitive effect against the tracer by the presence of unlabelled somatostatin
Different affinity profile of the analogue for the various sstr subtypes and variability of expression of the receptors: subtypes of receptors expressed and density of sstr expression over the tumour cells may influence the tumour detectability
Liver metastases may be taking up the tracer with a similar degree as the normal liver

Imaging results

The sensitivity of SRS by means of ^{111}In -DTPA-octreotide scan is well documented in the diagnosis pathways for NETs but may vary depending on the type of tumour studied and on the site of the disease. As already said one of the major issues with the study of NETs is the heterogeneity of these neoplasms.

Somatostatin receptor-positive tumours include pituitary adenomas, gastroenteropancreatic (GEP) tumours, carcinoids, small cell lung cancers (SCLC), paragangliomas, pheochromocytomas and neuroblastomas and medullary thyroid cancer. Some breast cancers have been reported and among the non-oncological diseases sarcoidosis can show positive results at SRS [10, 53]

Kwekkeboom et al. [54] recently listed the SRS results in tumours and other diseases (Table 4).

Pituitary

Somatostatin receptors have been demonstrated in vitro in pituitary adenomas and positive results at SRS have been reported in patients affected by functioning and non-functioning pituitary adenomas, with the uptake higher in the first group of patients than in the second [55, 56]. However, in pituitary adenomas the role of SRS is still limited because of the physiological uptake in the same site that makes the diagnostic accuracy poorer [55].

GEP NETs and carcinoids

GEPs and carcinoids are relatively rare diseases with a wide range of clinical presentations and therefore require a complex diagnostic workup in which scintigraphic imaging plays a relevant role both alone or in association with other instrumental examinations (Fig. 3).

The use of SRS in GEPs is largely diffused and there is nowadays quite a consolidated experience about it, supported by clinical data on its accuracy which is higher than conventional imaging (i.e. CT or MRI) with a sensitivity of 80–100% in localizing the primary tumour and disease burden [57, 58]. The association with other imaging techniques may improve the already high accuracy [59].

In a multicentre trial in Europe Krenning et al. [60] studied the sensitivity of histologically or biochemically proven pancreatic NET. The results were promising with a detection rate of 100% for glucagonomas, 88% for VIPomas, 72% for gastrinomas, 82% for non-functioning islet cell tumours and 87% for carcinoids.

Carcinoid tumour detection has been reported at nearly a 100% sensitivity and many authors describe the detection of unknown and unexpected sites of diseases that were not found at other imaging modalities [61]. This is very helpful

Table 4 Sensitivity of somatostatin receptor imaging modified from Kwekkeboom et al. [54]

Disease	Reference
High sensitivity (detection rate >75%)	
Pituitary tumours	Kwekkeboom et al. (1999) [55]
Gastrinomas	de Kerviler et al. (1994) [92]; Gibril et al. (1999) [49]
Non-functioning pancreatic NETs	Krenning et al. (1993) [13]; Lebtahi et al. (1997) [93]
Functioning pancreatic NETS except insulinomas	Krenning et al. (1993) [13]; Lebtahi et al. (1997) [93]
Carcinoids	Kwekkeboom et al. (1993) [94]; Westlin et al. (1993) [61]; Ahlman et al. (1994) [95]; Kälkner et al. (1995) [96]
Small cell lung cancer	Kirsch et al. (1994) [72]; Kwekkeboom et al. (1994) [71]; Bombardieri et al. (1995) [69]; Reisinger et al. (1998) [68]
Paragangliomas	Kwekkeboom et al. (1993) [81]; Telischi et al. (2000) [97]; Duet et al. (2003) [98]
Meningiomas	Haldemann et al. (1995) [99]; Schmidt et al. (1998) [100]
Sarcoidosis and other granulomatous diseases	Vanhagen et al. (1994) [101]; Kwekkeboom et al. (1998) [102]
Graves' disease and Graves' ophthalmopathy	Postema et al. (1994) [103]; Krassas et al. (1995) [104]
Intermediate sensitivity (detection rate ranging between 40 and 75%)	
Insulinoma	Krenning et al. (1993) [13]; Zimmer et al. (1996) [62]; Schillaci et al. (2000) [63]; Vezzosi et al. (2005) [64]
Medullary thyroid carcinoma	Kwekkeboom et al. (1993) [84]; Tisell et al. (1997) [105]; Adams et al. (1998) [106]
Differentiated thyroid carcinoma, including Hürthle cell carcinoma	Postema et al. (1996) [107]; Haslinghuis et al. (2001) [108]
Breast cancer	van Eijck et al. (1994) [109]
Lymphoma (NHL, HL)	Lugtenburg et al. (2001) [110, 111]
Phaeochromocytoma	van der Harst et al. (2001) [112]
Astrocytoma	Schmidt et al. (1998) [100]

NHL non-Hodgkin's lymphoma,
HL Hodgkin's lymphoma

as it can modify the therapeutic strategy in the management of the patient.

The reported sensitivity for the diagnosis of insulinomas is lower and ranges between 20 and 60%. [13, 62–64]. The density of expression of somatostatin receptor is different in benign and malignant insulinomas and a higher number of ^{111}In -DTPA-pentetreotide-positive scans have been reported [65].

In patients affected by malignant insulinomas, uptake in the primary and in the metastatic foci has been demonstrated in several studies [64, 66], although the high sensitivity of endoscopic ultrasound [67] is well known and the real role of SRS in these tumours still needs further investigations.

Small cell lung cancers

SRS positivity has been demonstrated in lung cancers and shows high sensitivity particularly in SCLC. In the multicentre study by Reisinger et al. [68] including 4 centres and 100 patients, ^{111}In -pentetreotide scintigraphy yielded a sensitivity of 96% in diagnosing the primary

tumour. Overall sensitivity in detecting the metastases was 54%, being higher for regional metastases (60%) and lower for distant metastases (45%). In an article published by Bombardieri et al. [69] the primary tumours were detected with a sensitivity of 95% and the metastases with a sensitivity of 80%. For the detection of the primary, Krenning et al. [13], O'Byrne et al. [70] and Kwekkeboom et al. [71] reported a sensitivity of 100%; Kirsch et al. [72] reported a sensitivity of 96% and Maini et al. [73] a sensitivity of 87%. The average sensitivity in the reported literature in detecting metastases is 59%.

Despite the limits in the detection of distant metastases, SRS by means of ^{111}In -pentetreotide is a useful tool for the evaluation of the extent of disease and may upstage patients from a limited disease to an extensive disease status.

Leitha et al. [74] reported a sensitivity of 84% for the primary and 65% for the metastases in 20 patients affected by SCLC and evaluated with ^{123}I -Tyr³-octreotide.

$^{99\text{m}}\text{Tc}$ -depreotide approved specifically for lung nodules has been reported with a sensitivity of 97% and a specificity of 73% [75].




Fig. 3 Imaging with ^{111}In -pentetreotide of gastrinoma. Patient with gastrinoma metastases, showing multiple lesions to liver, lung and bone

Phaeochromocytomas and paragangliomas

Phaeochromocytoma is a neoplasm arising from the adrenal medulla occurring in 0.1% of patients with hypertension and is frequently associated with pallor, headaches and palpitations; 10% of phaeochromocytomas are bilateral, 10% are malignant and 10% arise outside the adrenals [76].

SRS sensitivity in phaeochromocytomas has been reported to be comparable to metaiodobenzylguanidine (MIBG) scintigraphy, ranging between 86 and 100% as in the papers by Tenenbaum et al. [77], Hoefnagel et al. [78] and Krenning et al. [13].

Paragangliomas, rare tumours most frequently originating from aorticosympathetic paraganglia including the organs of Zuckerkandl, produce catecholamine [79].

Telischi et al. [80] described ^{111}In -pentetreotide scintigraphy having a high accuracy in the study of paragangliomas (90%), a 94% sensitivity and a specificity of 75%.

Ten per cent of paragangliomas are metastatic and the visualization in SRS can be of additional value to conventional imaging [81].

Phaeochromocytomas and paragangliomas show SRS-positive imaging, though $^{131/123}\text{I}$ -MIBG imaging is usually preferred for the study of the adrenal region which can be difficult in SRS given the kidney excretion of the tracer and the consequent high background activity.

Medullary thyroid cancer

Different single photon emission radiopharmaceuticals have been proposed for the study of medullary thyroid carcinoma. $^{99\text{m}}\text{Tc(V)}$ -DMSA showed an overall sensitivity of 69% as opposed to the 29% overall sensitivity shown by ^{111}In -pentetreotide in a study by Kurtaran et al. [82], but according to other experiences, SRS has been compared to conventional imaging [83] and has been reported with a sensitivity of 50–70% for medullary thyroid carcinoma [84].

Summary of clinical applications

The use of SRS in NET, above all in GEPs and carcinoids, is well established worldwide for the management of the patients and, as in the aforementioned ^{111}In -pentetreotide EANM guidelines [38], suited to:

- Localize primary tumours and detect sites of metastatic disease (staging)
- Follow up patients with known disease to detect residual, recurrent or progressive disease (restaging)
- Monitor the effects of therapy (surgery, radiotherapy, chemotherapy or somatostatin analogue therapy)
- Select patients for PRRT
- Obtain prognostic parameters for the response to subsequent therapy

In NET a generally high level of ssr expression is expected, but the heterogeneity of their distribution can be responsible for some discrepancies in the clinical features and in the imaging results. However, the major advantages of the SRS seem to be the possibility of selecting patients for PRRT and providing prognostic information since the cellular differentiation is thought

Table 5 Results of SRS, CT and US in the detection of primary and metastatic lesions from GEP NETs in 131 patients modified from Chiti et al. [89]

	SRS	CT	US
Primary tumour			
Sensitivity (%)	62	43	36
Liver metastases			
Sensitivity (%)	90	78	88
Specificity (%)	97	93	95
Accuracy (%)	93	83	91
Other soft tissue lesions			
Sensitivity (%)	90	66	47
Specificity (%)	98	98	100
Accuracy (%)	95	83	67

to be associated with high levels of ssr expression on the cell surface [85].

Somatostatin receptor scintigraphy versus conventional imaging

The heterogeneity of NET has made clear the need for an integrated workup both in diagnosis and therapy: SRS has shown to be an accurate tool for diagnosis but of course conventional radiological imaging, including abdominal computed tomography (CT), ultrasound (US) and magnetic resonance imaging (MRI), is required as well.

There is not an extensive number of studies in which SSR and conventional imaging have been compared. Zimmer et al. [86] reported in metastasized NET of the upper gastrointestinal tract a 52% detection rate for SRS, 36% for CT and 24% for MRI.

In another study by Zimmer et al. [62] ten gastrinoma patients and ten insulinoma patients were studied with SSR and conventional imaging. In a lesion-based analysis SRS showed an 86% sensitivity, whereas the combination of CT, US and MRI showed a 29% sensitivity.

In a study by Shi et al. [87] SRS, CT and MRI were compared in a group of 48 patients with known or clinically suspected NET. The sensitivity shown for SSR was 95%. In a lesion-based analysis SRS showed a detection rate of 87%, CT of 44% and MRI of 43%.

In 80 patients affected by Zollinger-Ellison syndrome studied by Gibril et al. [88], SRS succeeded in localizing the primary gastrinoma in 56% of patients, with a higher sensitivity than any conventional study including angiography.

We report here the interesting data from Chiti et al. [89]: in 131 patients with known or suspected NET of the GEP tract, the primary was identified by SRS with a sensitivity of 62%. The sensitivity in the detection of the primary fell to 43% for CT and to 36% for US. Considering the liver metastases there was a comparable sensitivity for SRS and US (90 vs 88%) and a lower one for CT (78%) (Table 5).

In a smaller series of patients in a more recent study, Schilaci [90] described a lower sensitivity for conventional imaging as compared to SRS in the detection of NET of the GEP tract.

However, we believe there is a need to compare the new improved conventional imaging technique with SRS in a larger series of patients.

Future role and perspectives

The basic biochemistry and physiology of somatostatin and octreotide have been dealt with in a recent review [91]. It brings a series of challenges for both basic science

researchers as well as for imaging experts. The “romantic” past of having one ligand and one receptor has to be re-evaluated in view of the complexities of receptor molecules (homodimers, heterodimers and truncated forms). This form of “socializing” might lead to the development of bivalent tracers and even to multi-agent diagnostic and therapeutic approaches. These thoughts might be applicable in many other systems based on other peptides. New theoretical models based on fractal dynamics that describe receptor interactions and OMICS might turn out to be the language of times to come and those who take up the challenge will write new reviews on this topic. A “brave new world” should come.

Conflicts of interest None.

References

1. Brazeau P, Vale W, Burgus R, Ling N, Butcher M, Rivier J, et al. Hypothalamic polypeptide that inhibits the secretion of immunoreactive pituitary growth hormone. *Science* 1973;179:77–9.
2. Plewe G, Beyer J, Krause U, Neufeld M, del Pozo E. Long-acting and selective suppression of growth hormone secretion by somatostatin analogue SMS 201–995 in acromegaly. *Lancet* 1984;2:782–4.
3. Patel YC, Greenwood MT, Warszynska A, Panetta R, Srikant CB. All five cloned somatostatin receptors (hSSTR1–5) are functionally coupled to adenylyl cyclase. *Biochem Biophys Res Commun* 1994;198:605–12.
4. Pangerl T, Peck-Radosavljevic M, Kaserer K, Niederle B, Gangl A, Virgolini I. Somatostatin (SST) and VIP receptor subtype gene expression in human tumors. *Eur J Nucl Med* 1997;24:995A.
5. Reubi JC. Peptide receptors as molecular targets for cancer diagnosis and therapy. *Endocr Rev* 2003;24:389–427.
6. Rufini V, Calcagni ML, Baum RP. Imaging of neuroendocrine tumors. *Semin Nucl Med* 2006;36(3):228–47.
7. Krenning EP, Bakker WH, Breeman WAP, Koper JW, Kooij PP, Ausema L, et al. Localisation of endocrine-related tumours with radiolabeled analogue of somatostatin. *Lancet* 1989;1:242–5.
8. Virgolini I, Yang Q, Li S, Angelberger P, Neuhold N, Niederle B, et al. Cross-competition between vasoactive intestinal peptide and somatostatin for binding to tumor cell membrane receptors. *Cancer Res* 1994;54:690–700.
9. Reubi JC, Schaer JC, Waser B, Mengod G. Expression and localization of somatostatin receptor SSTR1, SSTR2, and SSTR3 messenger RNAs in primary human tumors using in situ hybridization. *Cancer Res* 1994;54:3455–9.
10. Volante M, Bozzalla-Cassione F, Papotti M. Somatostatin receptors and their interest in diagnostic pathology. *Endocr Pathol* 2004;15(4):275–92.
11. de Herder WW, Lamberts SW. Somatostatin and somatostatin analogues: diagnostic and therapeutic uses. *Curr Opin Oncol* 2002;14:53–7.
12. Lamberts SWJ, Bakker WH, Reubi JC, Krenning EP. Somatostatin-receptor imaging in the localization of endocrine tumors. *N Engl J Med* 1990;323:1246–9.
13. Krenning EP, Kwekkeboom DJ, Bakker WH, Breeman WA, Kooij PP, Oei HY, et al. Somatostatin receptor scintigraphy with [111In-DTPA-D-Phe1]- and [123I-Tyr3]-octreotide: the Rotterdam experience with more than 1000 patients. *Eur J Nucl Med* 1993;20:716–31.

14. Raynor K, Lucki I, Reisine T. Somatostatin receptors in the nucleus accumbens selectively mediate the stimulatory effect of somatostatin on locomotor activity in rats. *J Pharmacol Exp Ther* 1993;265:67–73.
15. Lamberts SW, Krenning EP, Reubi JC. The role of somatostatin and its analogs in the diagnosis and treatment of tumors. *Endocr Rev* 1991;12:450–82.
16. Hoyer D, Bell GI, Berelowitz M, Epelbaum J, Feniuk W, Humphrey PP, et al. Classification and nomenclature of somatostatin receptors. *Trends Pharmacol Sci* 1995;16:86–8.
17. Lamberts SWJ, van der Lely AJ, de Herder WW, Hofland LJ. Drug therapy: octreotide. *N Engl J Med* 1996;334:246–54.
18. Smith-Jones P, Bischof C, Leimer M, Gludovac D, Angelberger P, Pangerl T, et al. “MAURITIUS” a novel somatostatin analog for tumor diagnosis and therapy. *J Nucl Med* 1998;39:223A.
19. Krenning EP, Bakker WH, Kooij PPM, Breeman WAP, Oei HY, de Jong M, et al. Somatostatin receptor scintigraphy with indium-111-DTPA-D-Phe-1-octreotide in man: metabolism, dosimetry and comparison with iodine-123-Tyr-3-octreotide. *J Nucl Med* 1992;33:652–8.
20. Bangard M, Béhé M, Gohlke S, Otte R, Bender H, Maecke HR, et al. Detection of somatostatin receptor-positive tumours using the new 99mTc-tricine-HYNIC-D-Phe1-Tyr3-octreotide: first results in patients and comparison with 111In-DTPA-D-Phe1-octreotide. *Eur J Nucl Med* 2000;27(6):628–37.
21. Decristoforo C, Mather SJ, Cholewinski W, Donnemiller E, Riccabona G, Moncayo R. 99mTc-EDDA/HYNIC-TOC: a new 99mTc-labelled radiopharmaceutical for imaging somatostatin receptor-positive tumours; first clinical results and intra-patient comparison with 111In-labelled octreotide derivatives. *Eur J Nucl Med* 2000;27:1318–25.
22. Kainz H, Bale R, Donnemiller E, Gabriel M, Kovacs P, Decristoforo C, et al. Image fusion analysis of (99m)Tc-HYNIC-octreotide scintigraphy and CT/MRI in patients with thyroid-associated orbitopathy: the importance of the lacrimal gland. *Eur J Nucl Med Mol Imaging* 2003;30:1155–9.
23. Krenning EP, Kooij PP, Pauwels S, Breeman WA, Postema PT, De Herder WW, et al. Somatostatin receptor: scintigraphy and radionuclide therapy. *Digestion* 1996;57:57–61.
24. Wild D, Schmitt JS, Ginj M, Mäcke HR, Bernard BF, Krenning E, et al. DOTA-NOC, a high-affinity ligand of somatostatin receptor subtypes 2, 3 and 5 for labelling with various radiometals. *Eur J Nucl Med Mol Imaging* 2003;30:1338–47.
25. Reubi JC, Schär JC, Waser B, Wenger S, Heppeler A, Schmitt JS, et al. Affinity profiles for human somatostatin receptor subtypes SST1–SST5 of somatostatin radiotracers selected for scintigraphic and radiotherapeutic use. *Eur J Nucl Med* 2000;27:273–82.
26. Ginj M, Chen J, Walter MA, Eltschinger V, Reubi JC, Maecke HR. Preclinical evaluation of new and highly potent analogues of octreotide for predictive imaging and targeted radiotherapy. *Clin Cancer Res* 2005;11:1136–45.
27. De Jong M, Breeman WA, Bakker WH, Kooij PP, Bernard BF, Hofland LJ, et al. Comparison of (111)In-labeled somatostatin analogues for tumor scintigraphy and radionuclide therapy. *Cancer Res* 1998;58:437–41.
28. Stahl A, Meisetschläger G, Schottelius M, Bruus-Jensen K, Wolf I, Scheidhauer K. [123I]Mtr-TOCA, a radioiodinated and carboxylated analogue of octreotide: scintigraphic comparison with [111In]octreotide. *Eur J Nucl Med Mol Imaging* 2006;33(1):45–52.
29. Maina T, Nock B, Nikolopoulou A, Sotiriou P, Loudos G, Maintas D, et al. [99mTc]Demotate, a new 99mTc-based [Tyr3] octreotide analogue for the detection of somatostatin receptor-positive tumours: synthesis and preclinical results. *Eur J Nucl Med Mol Imaging* 2002;29(6):742–53.
30. Maina T, Nock BA, Cordopatis P, Bernard BF, Breeman WA, van Gameren A, et al. [(99m)Tc]Demotate 2 in the detection of sst(2)-positive tumours: a preclinical comparison with [(111)In]DOTA-tate. *Eur J Nucl Med Mol Imaging* 2006;33(7):831–40.
31. Decristoforo C, Maina T, Nock B, Gabriel M, Cordopatis P, Moncayo R. 99mTc-Demotate 1: first data in tumour patients—results of a pilot/phase I study. *Eur J Nucl Med Mol Imaging* 2003;30:1211–9.
32. Rogers BE, Parry JJ, Andrews R, Cordopatis P, Nock BA, Maina T. MicroPET imaging of gene transfer with a somatostatin receptor-based reporter gene and (94m)Tc-Demotate 1. *J Nucl Med* 2005;46:1889–97.
33. Gabriel M, Decristoforo C, Wöll E, Eisterer W, Nock B, Maina T, et al. [99mTc]Demotensin VI: biodistribution and initial clinical results in tumor patients of a pilot/phase I study. *Cancer Biother Radiopharm* 2011;26(5):557–63.
34. Virgolini I, Britton K, Buscombe J, Moncayo R, Paganelli G, Riva P. 111In- and 90Y-DOTA-lanreotide: results and implications of the MAURITIUS trial. *Semin Nucl Med* 2002;32:148–55.
35. Menda Y, Kahn D. Somatostatin receptor imaging of non-small cell lung cancer with 99mTc depreotide. *Semin Nucl Med* 2002;32:92–6.
36. Lebtahi R, Le Cloirec J, Houzard C, Daou D, Sobhani I, Sassolas G, et al. Detection of neuroendocrine tumors: (99m)Tc-P829 scintigraphy compared with (111)In-pentetreotide scintigraphy. *J Nucl Med* 2002;43:889–95.
37. Bogatzky J, Gohlke S, Bender H, Sartor J, Knapp FF, Biersack HJ. Receptor binding studies with rhenium-RC160, a directly-labeled somatostatin analog, with membranes of the colon cancer cell line HT 29, CHO cells and ovary carcinoma cell line EFO-21. *Eur J Nucl Med* 1997;24:1059A.
38. Bombardieri E, Ambrosini V, Aktolun C, Baum RP, Bischof-Delaloye A, Del Vecchio S, et al. 111In-pentetreotide scintigraphy: procedure guidelines for tumour imaging. *Eur J Nucl Med Mol Imaging* 2010;37:1441–8.
39. ICRP Publication 80. Radiation dose to patients from radiopharmaceuticals. *Ann ICRP* 1998;28:3. Oxford: Pergamon.
40. ICRP Publication 62. Radiological protection in biomedical research. *Ann ICRP* 1991;22:3. Oxford: Pergamon.
41. ICRP Publication 53. Radiation dose to patients from radiopharmaceuticals. *Ann ICRP* 1987;18:1–4. Oxford: Pergamon.
42. ICRP Publication 106. Radiation dose to patients from radiopharmaceuticals. Addendum 3 to ICRP Publication 53. Approved by the Commission in October 2007. *Ann ICRP* 2008;38(1–2):1–197.
43. Steffen IG, Mehl S, Heuck F, Elgeti F, Furth C, Amthauer H, et al. Attenuation correction of somatostatin receptor SPECT by integrated low-dose CT: is there an impact on sensitivity? *Clin Nucl Med* 2009;34(12):869–73.
44. Hillel PG, van Beek EJ, Taylor C, Lorenz E, Bax ND, Prakash V, et al. The clinical impact of a combined gamma camera/CT imaging system on somatostatin receptor imaging of neuroendocrine tumours. *Clin Radiol* 2006;61(7):579–87.
45. Ruf J, Steffen I, Mehl S, Rosner C, Denecke T, Pape UF, et al. Influence of attenuation correction by integrated low-dose CT on somatostatin receptor SPECT. *Nucl Med Commun* 2007;28(10):782–8.
46. Gabriel M, Hausler F, Bale R, Moncayo R, Decristoforo C, Kovacs P, et al. Image fusion analysis of (99m)Tc-HYNIC-Tyr (3)-octreotide SPECT and diagnostic CT using an immobilisation device with external markers in patients with endocrine tumours. *Eur J Nucl Med Mol Imaging* 2005;32:1440–51.
47. Jacobsson H, Bremmer S, Larsson SA. Visualisation of the normal adrenals at SPET examination with 111In-pentetreotide. *Eur J Nucl Med Mol Imaging* 2003;30:1169–72.
48. Kwekkeboom DJ, Krenning EP. Somatostatin receptor imaging. *Semin Nucl Med* 2002;32(2):84–91.
49. Gibril E, Reynolds JC, Chen CC, Yu F, Goebel SU, Serrano J, et al. Specificity of somatostatin receptor scintigraphy: a prospective

- study and effects of false-positive localizations on management in patients with gastrinomas. *J Nucl Med* 1999;40:539–53.
50. Krausz Y, Keidar Z, Kogan I, Even-Sapir E, Bar-Shalom R, Engel A, et al. SPECT/CT hybrid imaging with ¹¹¹In-pentetreotide in assessment of neuroendocrine tumours. *Clin Endocrinol (Oxf)* 2003;59:565–73.
51. Moreira AP, Duarte LH, Vieira F, João F, Lima JP. Value of SPET/CT image fusion in the assessment of neuroendocrine tumours with ¹¹¹In-pentetreotide scintigraphy. *Rev Esp Med Nucl* 2005;24:14–8.
52. Perri M, Erba P, Volterrani D, Lazzeri E, Boni G, Grosso M, et al. Octreo-SPECT/CT imaging for accurate detection and localization of suspected neuroendocrine tumors. *Q J Nucl Med Mol Imaging* 2008;52:323–33.
53. Briganti V, Sestini R, Orlando C, Bernini G, La Caya G, Tamburini A, et al. Imaging of somatostatin receptors by indium-111-pentetreotide correlates with quantitative determination of somatostatin receptor type 2 gene expression in neuroblastoma tumors. *Clin Cancer Res* 1997;3:2385–91.
54. Kwekkeboom DJ, Kam BL, van Essen M, Teunissen JJM, van Eijck CH, Valkema R, et al. Somatostatin-receptor-based imaging and therapy of gastroenteropancreatic neuroendocrine tumors. *Endocr Relat Cancer* 2010;17:R53–73.
55. Kwekkeboom DJ, de Herder WW, Krenning EP. Receptor imaging in the diagnosis and treatment of pituitary tumors. *J Endocrinol Invest* 1999;22:80–8.
56. Duet M, Mundler O, Azjenberg C, Berolatti B, Chedin P, Duranteau L, et al. Somatostatin receptor imaging in non-functioning pituitary adenomas: value of an uptake index. *Eur J Nucl Med* 1994;21:647–50.
57. Chiti A, Briganti V, Fanti S, Monetti N, Masi R, Bombardieri E. Results and potential of somatostatin receptor imaging in neuroendocrine gastroenteropancreatic tumours. *Q J Nucl Med* 2000;44:42–9.
58. Briganti V, Matteini M, Ferri P, Vaggelli L, Castagnoli A, Pieroni C. Octreoscan SPET evaluation in the diagnosis of pancreas neuroendocrine tumors. *Cancer Biother Radiopharm* 2001;16:515–24.
59. Bombardieri E, Seregini E, Villano C, Chiti A, Bajetta E. Position of nuclear medicine techniques in the diagnostic work-up of neuroendocrine tumors. *Q J Nucl Med Mol Imaging* 2004;48:150–63.
60. Krenning EP, Kwekkeboom DJ, Pauwels S, Kvols LK, Reubi JC. Somatostatin receptor scintigraphy. In: Freeman LM, editor. *Nuclear medicine annual*. New York: Lippincott Williams & Wilkins; 1995. p. 1–50.
61. Westlin J, Janson ET, Arnberg H, Ahlström H, Öberg K, Nilsson S. Somatostatin receptor scintigraphy of carcinoid tumours using the [¹¹¹In-DTPA-D-Phe¹]-octreotide. *Acta Oncol* 1993;32:783–6.
62. Zimmer T, Stölzel U, Bäder M, Koppenhagen K, Hamm B, Buhr H, et al. Endoscopic ultrasonography and somatostatin receptor scintigraphy in the preoperative localisation of insulinomas and gastrinomas. *Gut* 1996;39:562–8.
63. Schillaci O, Massa R, Scopinaro F. ¹¹¹In-pentetreotide scintigraphy in the detection of insulinomas: importance of SPECT imaging. *J Nucl Med* 2000;41:459–62.
64. Vezzosi D, Bennet A, Rochaix P, Courbon F, Selves J, Pradere B, et al. Octreotide in insulinoma patients: efficacy on hypoglycemia, relationships with OctreoScan scintigraphy and immunostaining with anti-sst2A and anti-sst5 antibodies. *Eur J Endocrinol* 2005;152:757–67.
65. Bertherat J, Tenenbaum F, Perlemoine K, Videau C, Alberini JL, Richard B, et al. Somatostatin receptors 2 and 5 are the major somatostatin receptors in insulinomas: an in vivo and in vitro study. *J Clin Endocrinol Metab* 2003;88:5353–60.
66. Baldelli R, Ettorre G, Vennarecci G, Pasimeni G, Carboni F, Lorusso R, et al. Malignant insulinoma presenting as metastatic liver tumor. Case report and review of the literature. *J Exp Clin Cancer Res* 2007;26:603–7.
67. de Geus-Oei LF, Oei HY, Hennemann G, Krenning EP. Sensitivity of ¹²³I whole-body scan and thyroglobulin in the detection of metastases or recurrent differentiated thyroid cancer. *Eur J Nucl Med Mol Imaging* 2002;29:768–74.
68. Reisinger I, Bohuslavitzki KH, Brenner W, Braune S, Ditttrich I, Geide A, et al. Somatostatin receptor scintigraphy in small-cell lung cancer: results of a multicenter study. *J Nucl Med* 1998;39:224–7.
69. Bombardieri E, Crippa F, Cataldo I, Chiti A, Seregini E, Soresi E, et al. Somatostatin receptor imaging of small cell lung cancer (SCLC) by means of ¹¹¹In-DTPA octreotide scintigraphy. *Eur J Cancer* 1995;31A:184–8.
70. O'Byrne KJ, Ennis JT, Freyne PJ, Clancy LJ, Prichard JS, Carney DN. Scintigraphic imaging of small-cell lung cancer with [¹¹¹In] pentetreotide, a radiolabelled somatostatin analogue. *Br J Cancer* 1994;69:762–6.
71. Kwekkeboom DJ, Kho GS, Lamberts SW, Reubi JC, Laissue JA, Krenning EP. The value of octreotide scintigraphy in patients with lung cancer. *Eur J Nucl Med* 1994;21:1106–13.
72. Kirsch CM, von Pawel J, Grau I, Tatsch K. Indium-111 pentetreotide in the diagnostic work-up of patients with bronchogenic carcinoma. *Eur J Nucl Med* 1994;21:1318–25.
73. Maini CL, Tofani A, Ventura I, Pigorini F, Sciuto R, Semprebene A, et al. Somatostatin receptor imaging in small cell lung cancer using ¹¹¹In-DTPA-octreotide: a preliminary study. *Nucl Med Commun* 1993;14:962–8.
74. Leitha T, Meghdadi S, Studnicka M, Wolzt M, Marosi C, Angelberger P, et al. The role of iodine-123-Tyr-3-octreotide scintigraphy in the staging of small-cell lung cancer. *J Nucl Med* 1993;34:1397–402.
75. Blum J, Handmaker H, Lister-James J, Rinne N. A multicenter trial with a somatostatin analog (^{99m}Tc depreotide) in the evaluation of solitary pulmonary nodules. *Chest* 2000;117:1232–8.
76. Cryer PE. The adrenal medullae. In: Wyngaarden JB, Smith LH, Bennet JC, editors. *Cecil textbook of medicine*. 19th ed. Philadelphia: Saunders; 1992. p. 1390–4.
77. Tenenbaum F, Lumbroso J, Schlumberger M, Mure A, Plouin PF, Caillou B, et al. Comparison of radiolabeled octreotide and metaiodobenzylguanidine (MIBG) scintigraphy in malignant pheochromocytoma. *J Nucl Med* 1995;36:1–6.
78. Hoefnagel CA, Lewington VJ. MIBG therapy. In: Ell PJ, Gambhir SS, editors. *Nuclear medicine in clinical diagnosis and treatment*, Vol. 1. New York: Churchill Livingstone; 2004. p. 445–557.
79. DeLellis RA. Multiple endocrine neoplasia syndromes revisited. Clinical, morphologic, and molecular features. *Lab Invest* 1995;72:494–502.
80. Telischi FF, Bustillo A, Whiteman ML, Serafini AN, Reisberg MJ, Gomez-Marin O, et al. Octreotide scintigraphy for the detection of paragangliomas. *Otolaryngol Head Neck Surg* 2000;122(3):358–62.
81. Kwekkeboom DJ, van Urk H, Pauw KH, Lamberts SW, Kooij PP, Hoogma RP, et al. Octreotide scintigraphy for the detection of paragangliomas. *J Nucl Med* 1993;34:873–8.
82. Kurtaran A, Leimer M, Kaserer K, Yang Q, Angelberger P, Niederle B, et al. Combined use of ¹¹¹In-DTPA-D-Phe-1-octreotide (OCT) and ¹²³I-vasoactive intestinal peptide (VIP) in the localization diagnosis of medullary thyroid carcinoma (MTC). *Nucl Med Biol* 1996;23(4):503–7.
83. Baudin E, Lumbroso J, Schlumberger M, Leclere J, Giammarile F, Gardet P, et al. Comparison of octreotide scintigraphy and

- conventional imaging in medullary thyroid carcinoma. *J Nucl Med* 1996;37:912–6.
84. Kwekkeboom DJ, Reubi JC, Lamberts SWJ, Bruining HA, Mulder AH, Oei HY, et al. In vivo somatostatin receptor imaging in medullary thyroid carcinoma. *J Clin Endocrinol Metab* 1993;76:1413–7.
85. Asnacios A, Courbon F, Rochaix P, Bauvin E, Cances-Lauwers V, Susini C, et al. Indium-111-pentetreotide scintigraphy and somatostatin receptor subtype 2 expression: new prognostic factors for malignant well-differentiated endocrine tumors. *J Clin Oncol* 2008;26(6):963–70.
86. Zimmer T, Ziegler K, Bäder M, Fett U, Hamm B, Riecken O, et al. Localisation of neuroendocrine tumours of the upper gastrointestinal tract. *Gut* 1994;35:471–5.
87. Shi W, Johnston CF, Buchanan KD, Ferguson WR, Laird JD, Crothers JG, et al. Localization of neuroendocrine tumours with [¹¹¹In]DTPA-octreotide scintigraphy (Octreoscan): a comparative study with CT and MR imaging. *QJM* 1998;91(4):295–301.
88. Gibril F, Reynolds JC, Doppman JL, Chen CC, Venzon DJ, Termanini B, et al. Somatostatin receptor scintigraphy: its sensitivity compared with that of other imaging methods in detecting primary and metastatic gastrinomas. A prospective study. *Ann Intern Med* 1996;125:26–34.
89. Chiti A, Fanti S, Savelli G, Romeo A, Bellanova B, Rodari M, et al. Comparison of somatostatin receptor imaging, computed tomography and ultrasound in the clinical management of neuroendocrine gastro-entero-pancreatic tumours. *Eur J Nucl Med* 1998;25:1396–403.
90. Schillaci O. Functional–anatomical image fusion in neuroendocrine tumors. *Cancer Biother Radiopharm* 2004;19:129–34.
91. Moncayo R. Reflections on the theory of “silver bullet” octreotide tracers: implications for ligand–receptor interactions in the age of peptides, heterodimers, receptor mosaics, truncated receptors, and multifractal analysis. *EJNMMI Res* 2011;1:9.
92. de Kerviler E, Cadiot G, Lebtahi R, Faraggi M, Le Guludec D, Mignon M. Somatostatin receptor scintigraphy in forty-eight patients with the Zollinger–Ellison syndrome. *GRESZE: Groupe d'Etude du Syndrome de Zollinger–Ellison*. *Eur J Nucl Med* 1994;21:1191–7.
93. Lebtahi R, Cadiot G, Sarda L, Daou D, Faraggi M, Petegnief Y, et al. Clinical impact of somatostatin receptor scintigraphy in the management of patients with neuroendocrine gastroenteropancreatic tumors. *J Nucl Med* 1997;38:853–8.
94. Kwekkeboom DJ, Krenning EP, Bakker WH, Oei HY, Kooij PPM, Lamberts SWJ. Somatostatin analogue scintigraphy in carcinoid tumours. *Eur J Nucl Med* 1993;20:283–92.
95. Ahlman H, Wängberg B, Tisell LE, Nilsson O, Fjälling M, Forssell-Aronsson E. Clinical efficacy of octreotide scintigraphy in patients with midgut carcinoid tumours and evaluation of intraoperative scintillation detection. *Br J Surg* 1994;81:1144–9.
96. Kälkner KM, Janson ET, Nilsson S, Carlsson S, Oberg K, Westlin JE. Somatostatin receptor scintigraphy in patients with carcinoid tumors: comparison between radioligand uptake and tumor markers. *Cancer Res* 1995;55 Suppl 23:5801s–4s.
97. Telischi FF, Bustillo A, Whiteman ML, Serafini AN, Reisberg MJ, Gomez-Marin O, et al. Octreotide scintigraphy for the detection of paragangliomas. *Otolaryngol Head Neck Surg* 2000;122:358–62.
98. Duet M, Sauvaget E, Pételle B, Rizzo N, Guichard JP, Wassef M, et al. Clinical impact of somatostatin receptor scintigraphy in the management of paragangliomas of the head and neck. *J Nucl Med* 2003;44:1767–74.
99. Haldemann AR, Rösler H, Barth A, Waser B, Geiger L, Godoy N, et al. Somatostatin receptor scintigraphy in central nervous system tumors: role of blood–brain barrier permeability. *J Nucl Med* 1995;36:403–10.
100. Schmidt M, Scheidhauer K, Luyken C, Voth E, Hildebrandt G, Klug N, et al. Somatostatin receptor imaging in intracranial tumours. *Eur J Nucl Med* 1998;25:675–86.
101. Vanhagen PM, Krenning EP, Reubi JC, Kwekkeboom DJ, Bakker WH, Mulder AH, et al. Somatostatin analogue scintigraphy in granulomatous diseases. *Eur J Nucl Med* 1994;21:497–502.
102. Kwekkeboom DJ, Krenning EP, Kho GS, Breeman WA, Van Hagen PM. Somatostatin receptor imaging in patients with sarcoidosis. *Eur J Nucl Med* 1998;25:1284–92.
103. Postema PTE, Krenning EP, Wijngaarde R, Kooy PP, Oei HY, van den Bosch WA, et al. [¹¹¹In-DTPA-D-Phe1] octreotide scintigraphy in thyroidal and orbital Graves' disease: a parameter for disease activity? *J Clin Endocrinol Metab* 1994;79:1845–51.
104. Krassas GE, Dumas A, Pontikides N, Kaltsas T. Somatostatin receptor scintigraphy and octreotide treatment in patients with thyroid eye disease. *Clin Endocrinol (Oxf)* 1995;42:571–80.
105. Tisell LE, Ahlman H, Wängberg B, Hansson G, Mölne J, Nilsson O, et al. Somatostatin receptor scintigraphy in medullary thyroid carcinoma. *Br J Surg* 1997;84:543–7.
106. Adams S, Baum RP, Hertel A, Schumm-Draeger PM, Usadel KH, Hör G. Comparison of metabolic and receptor imaging in recurrent medullary thyroid carcinoma with histopathological findings. *Eur J Nucl Med* 1998;25:1277–83.
107. Postema PTE, De Herder WW, Reubi JC, Oei HY, Kwekkeboom DJ, Bruining HJ, et al. Somatostatin receptor scintigraphy in non-medullary thyroid cancer. *Digestion* 1996;57:36–7.
108. Haslinghuis LM, Krenning EP, de Herder WW, Kwekkeboom DJ. Somatostatin receptor scintigraphy in the follow-up of patients with differentiated thyroid cancer. *J Endocrinol Invest* 2001;24:415–22.
109. van Eijck CH, Krenning EP, Bootsma A, Oei HY, van Pel R, Lindemans J, et al. Somatostatin–receptor scintigraphy in primary breast cancer. *Lancet* 1994;343:640–3.
110. Lugtenburg PJ, Löwenberg B, Valkema R, Oei HY, Lamberts SW, Eijkemans MJ, et al. Somatostatin receptor scintigraphy in the initial staging of low-grade non-Hodgkin's lymphomas. *J Nucl Med* 2001;42:222–9.
111. Lugtenburg PJ, Krenning EP, Valkema R, Oei HY, Lamberts SW, Eijkemans MJ, et al. Somatostatin receptor scintigraphy useful in stage I–II Hodgkin's disease: more extended disease identified. *Br J Haematol* 2001;112:936–44.
112. van der Harst E, de Herder WW, Bruining HA, Bonjer HJ, de Krijger RR, Lamberts SW, et al. [(123)I]metaiodobenzylguanidine and [(111)In]octreotide uptake in benign and malignant pheochromocytomas. *J Clin Endocrinol Metab* 2001;86:685–93.