

Comparison of molecular markers to detect fresh sewage in environmental waters

W. Ahmed ^{*a,b,**}, A. Goonetilleke ^{*b*}, D. Powell ^{*c*}, K. Chauhan ^{*b*}, T. Gardner ^{*a,b*}

^a Department of Natural Resources and Water, 80 Meiers Road, Indooroopilly, Brisbane 4068, Australia ^b School of Urban Development, Queensland University of Technology, GPO Box 2434, Brisbane 4001, Australia ^c Faculty of Science, Health and Education, University of the Sunshine Coast, Maroochydore DC, 4558, Qld, Australia

ARTICLE INFO

Article history: Received 2 March 2009 Received in revised form 26 July 2009 Accepted 17 September 2009 Published online 22 September 2009

Keywords: Microbial source tracking Sewage pollution Molecular markers Enteric viruses Water quality

ABSTRACT

Human-specific Bacteroides HF183 (HS-HF183), human-specific Enterococci faecium esp (HS-esp), human-specific adenoviruses (HS-AVs) and human-specific polyomaviruses (HS-PVs) assays were evaluated in freshwater, seawater and distilled water to detect fresh sewage. The sewage spiked water samples were also tested for the concentrations of traditional fecal indicators (i.e., Escherichia coli, enterococci and Clostridium perfringens) and enteric viruses such as enteroviruses (EVs), sapoviruses (SVs), and torquetenoviruses (TVs). The overall host-specificity of the HS-HF183 marker to differentiate between humans and other animals was 98%. However, the HS-esp, HS-AVs and HS-PVs showed 100% hostspecificity. All the human-specific markers showed >97% sensitivity to detect human fecal pollution. E. coli, enterococci and, C. perfringens were detected up to dilutions of sewage 10⁻⁵, 10^{-4} and 10^{-3} respectively. HS-*esp*, HS-AVs, HS-PVs, SVs and TVs were detected up to dilution of sewage 10^{-4} whilst EVs were detected up to dilution 10^{-5} . The ability of the HS-HF183 marker to detect fresh sewage was 3-4 orders of magnitude higher than that of the HS-esp and viral markers. The ability to detect fresh sewage in freshwater, seawater and distilled water matrices was similar for human-specific bacterial and viral marker. Based on our data, it appears that human-specific molecular markers are sensitive measures of fresh sewage pollution, and the HS-HF183 marker appears to be the most sensitive among these markers in terms of detecting fresh sewage. However, the presence of the HS-HF183 marker in environmental waters may not necessarily indicate the presence of enteric viruses due to their high abundance in sewage compared to enteric viruses. More research is required on the persistency of these markers in environmental water samples in relation to traditional fecal indicators and enteric pathogens.

 $\ensuremath{\textcircled{}^\circ}$ 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Fecal pollution is one of the major concerns in relation to water bodies used for drinking water supply, recreational activities and harvesting seafood due to likely exposure to a wide array of pathogenic bacteria, protozoa and viruses (Hörman et al., 2004; Fong and Lipp, 2005). Various sources such as agricultural run-off, wild animals, combined sewer overflows (CSOs), sewage treatment plants (STPs), defective on-site wastewater treatment systems and industrial wastewater outlets are known to be potential sources of such pollution. The microbiological quality of water is generally assessed by enumerating fecal indicator bacteria such as *Escherichia coli* and enterococci which are commonly found in

E-mail address: shuhat@yahoo.com (W. Ahmed).

^{*} Corresponding author at: Department of Natural Resources and Water, 80 Meiers Road, Indooroopilly, Brisbane 4068, Australia. Tel.: +61 7 3896 9623.

^{0043-1354/\$ –} see front matter \circledast 2009 Elsevier Ltd. All rights reserved. doi:10.1016/j.watres.2009.09.047

the feces of warm-blooded animals including humans (USEPA, 2000). The presence of these indicators in water bodies generally points to fecal pollution and potential public health risks. The identification of indicator bacteria from major polluting source(s) is vitally important in order to implement appropriate mitigation strategies to minimise fecal pollution and associated public health risks (Scott et al., 2002). However, the assignment of indicator bacteria to human and animal sources in environmental waters is difficult due to their cosmopolitan nature (Field and Samadpour, 2007). In addition, environmental waters can be impacted by multiple sources of fecal pollution making it extremely difficult to implement a robust management plan without understanding the potential sources of pollution.

Over the last decade, microbial source tracking (MST) techniques have been developed to distinguish human from animal fecal pollution. The underlying assumption of MST is that the host-specificity of microorganisms is influenced by selective pressure in the host animal gut (Wiggins, 1996). The majority of the early MST methods are library-dependent which require the development of a collection of E. coli or enterococci isolates from suspected sources using various phenotypic and genotypic methods. Phenotypic or genotypic patterns of target strains are then compared to the library to identify their likely sources (Scott et al., 2002). There are several significant limitations in library-dependent methods which have been widely reported in the research literature such as: (1) a large representative library is required for successful field application. The development of such a library is laborious, and usually costly when using phenotypic and genotypic methods (i.e., PFGE and carbon source utilization) (Field and Samadpour, 2007); (2) commonly used fecal indicator bacteria (E. coli and/or enterococci) lack host-specificity (Gordon et al., 2002); (3) a library consisting of a small number of isolates cannot be readily used in multiple catchments, and therefore development of a separate library may be required for each catchment of interest (Ahmed et al., 2006; Hartel et al., 2002; Scott et al., 2003), and (4) library-dependent methods may yield both high false positive and negative results (Harwood et al., 2003; Moore et al., 2005).

Certain limitations of library-dependent methods could be overcome by using library-independent methods. These methods rely on detecting host-specific molecular markers in a given environmental sample using PCR assays. These methods are rapid and have shown to have higher specificities in a method comparison study (Griffith et al., 2003). The most commonly used markers for MST can be categorised into three groups: (1) anaerobic bacterial markers (i.e., hostspecific Bacteroides PCR) (Bernhard and Field, 2000), (2) bacterial toxin markers (i.e., E. faecium esp and E. coli toxin gene markers) (Scott et al., 2005; Khatib et al., 2002), and (3) viral markers (i.e., host-specific adenoviruses and polyomaviruses) (Fong and Lipp, 2005; McQuaig et al., 2006). Several studies have reported high host specificities of these markers which makes them suitable to distinguish between sources of fecal pollution (Ahmed et al., 2008a; Seurinck et al., 2005; Bernhard and Field, 2000; Reischer et al., 2006; Scott et al., 2005; Khatib et al., 2002).

Library-independent methods also have potential limitations, including the detection of certain markers in a small number of non-target samples (Gourmelon et al., 2007; Carson et al., 2005; Gawler et al., 2007; Whitman et al., 2007). Another limitation of these markers is that they are not present in the feces of all individuals, and the concentrations may vary from one DNA target to another (Field and Samadpour, 2007). For example, the concentration of human-specific Bacteroides markers in sewage samples could be 4-5 orders of magnitude higher than human-specific viral or toxin gene markers. Moreover, little is known regarding the persistency of these markers in environmental waters. In addition, the correlation between some of these markers with traditional fecal indicators and pathogens is not well documented. The absence of a particular marker in environmental waters does not completely rule out the presence of fecal pollution from that particular source. A general consensus is that multiple markers should be used (where possible) to obtain accurate and confirmatory results. To-date, only a few studies have used multiple host-specific markers to identify the sources of fecal pollution in environmental waters (Ahmed et al., 2007; Gourmelon et al., 2007; McQuaig et al., 2006). These markers appear to be promising in identifying the sources of fecal pollution. However, more research is required prior to their application for routine monitoring of water quality. A recent review paper highlighted the various research gaps that need to be addressed for library-independent methods (Santo Domingo et al., 2007).

The aim of this study was to evaluate the human-specific Bacteroides HF183 (HS-HF183), human-specific E. faecium esp (HS-esp), human-specific adenoviruses (HS-AVs), and humanspecific polyomaviruses (HS-PVs) markers to detect the smallest amount of fresh sewage pollution in sewage spiked freshwater, seawater and distilled water samples using realtime PCR assays. Furthermore, the sewage spiked water samples were also tested for the concentrations of fecal indicators such as E. coli, enterococci and Clostridium perfringens. In addition, real-time PCR assays were also used to detect enteric viruses such as emerging enteroviruses (EVs), sapoviruses (SVs), and torquetenoviruses (TVs) for sewage spiked water samples. These enteric viruses are excreted in extremely high numbers in the feces of infected individuals and can cause mild to severe gastroenteritis in humans. Humans could be exposed to enteric viruses by using contaminated waters for shellfish harvesting or recreation, or as a source of drinking waters. Fecal indicators' concentrations and the ability of each human-specific marker to detect fresh sewage were used to obtain a better understanding of which fecal indicators and human-specific marker(s) could potentially indicate the presence of enteric viruses in environmental waters polluted with fresh sewage.

2. Materials and methods

2.1. Host-specificity and sensitivity of human-specific markers

Host-specificity and sensitivity are commonly used parameters for human-specific markers. The specificity of a marker is the proportion of negative-control samples in which the marker is detected and the sensitivity of a marker is the proportion of positive control samples in which the marker is detected. To determine the host-specificity and sensitivity of the human-specific markers, 82 fecal samples were collected from six host groups. Approximately 10 ml of sample from each human fecal source (n = 32) was collected from influent entering sewage treatment plants (STPs). Cattle (n = 10), pigs (n = 10), and sheep (n = 10) samples (i.e., individual and composite animal wastewater samples) were collected from an abattoir in Killarney, Brisbane. Dog fecal samples (n = 10)were collected from two dog parks. Duck fecal samples (n = 10) were collected from Brisbane City Botanical Gardens where a large number of ducks roam. A fresh fecal sample (approximately 0.5–1.0 g) was collected from each individual animal (n = 50) with sterile swabs and inserted into a sterile container, transported on ice to the laboratory, stored at 4 °C, and processed within 6 h. For DNA extraction three different methods were used according to the previously published methods (Ahmed et al., 2008a,b; Haramoto et al., 2005). Approximately 180–200 mg of fresh animal feces was used for DNA extraction using each method separately.

2.2. Testing for real-time PCR inhibitors

Fecal and STP samples contain numerous organic and inorganic substances with the potential to inhibit PCR analyses (Wilson, 1997). An experiment was conducted to determine the potential presence of inhibitory substances in DNA extracted from animal fecal and STP samples for the HS-HF183, HS-*esp* and HS-AVs and HS-PVs assays. DNA was extracted from 1 L of ultrapure DNase- and RNase-free sterile distilled water (Invitrogen) after concentrating the sample. A representative number of pooled animal fecal samples (n = 5) and STP samples (n = 5) were spiked with 10³ gene copies of the HS-HF183 and cattle-specific CF128 (CS-CF128) markers (Bernhard and Field, 2000). The threshold cycle (C_T) values of these spiked DNA samples were compared to those of the DNA samples from distilled water spiked with the same concentration of the HS-HF183 and CS-CF128 markers.

2.3. Sample preparation

To determine the ability of each human-specific marker to detect fresh sewage, sewage samples were collected from the primary influent of a STP. Sewage samples were suspended in freshwater (n = 3), seawater (n = 3), and distilled water (n = 3) at a ratio 1:1 (62.5 ml water:62.5 ml fresh sewage), and a serial dilution ($10^{-1}-10^{-10}$) was made for each sample type. Environmental water samples were autoclaved before spiking, and exposed under UV light for 1 h to minimise background target DNA that could be present due to fecal pollution. The numbers of *E. coli*, enterococci, and *C. perfringens* were enumerated using membrane filtration method for each dilution. DNA and RNA extraction was performed for each dilution for the real-time PCR assays of the human-specific markers and enteric viruses.

2.4. Isolation and enumeration of fecal indicators

The membrane filtration method was used to process the sewage spiked water samples for *E. coli*, enterococci, and *C.*

perfringens enumeration. Sample serial dilutions were made and filtered through 0.45-µm pore size (47-mm diameter) nitrocellulose membranes (Advantec, Tokyo, Japan), and placed on modified membrane-thermotolerant E. coli agar (modified mTEC agar) (Difco, Detroit, MI, USA), membrane-Enterococcus indoxyl-D-glucoside (mEI) agar (Difco), and oleandomycin-polymyxin-sulfadiazine perfringens (OPSP) agar with supplement for the isolation of E. coli, enterococci, and C. perfringens, respectively. For the isolation of C. perfringens spores, water samples were heated at 60 °C for 30 min before filtration. Modified mTEC agar plates were incubated at 35 °C for 2 h to recover stressed cells, followed by incubation at 44 °C for 22 h (USEPA, 2002), and mEI agar plates were incubated at 41 °C for 48 h (USEPA, 1997). The OPSP agar plates were overlaid with 15 ml of molten OPSP agar before incubation. OPSP agar plates (for C. perfringens) were incubated anaerobically at 44 °C for 24 h. A confirmatory test for C. perfringens was performed according to the method described previously (Wohlsen et al., 2006).

2.5. Specificity of the real-time PCR primers

Previously published primers were used in this study. The primer sequence and annealing temperature for corresponding targets are shown in Table 1. Primer specificity was checked by searching for similar sequences in microbial genomes using the Basic Local Alignment Search Tool (BLAST) software (http://www.ncbi.nlm.nih.gov/BLAST/).

2.6. DNA and RNA extraction

DNA (i.e., HS-HF183) was extracted for each dilution according to the previously published method (Bernhard and Field, 2000). In brief, 125 ml of each sewage spiked water sample was filtered through 0.45 μ m pore size membranes (Advantec, Tokyo, Japan). The filters were transferred in sterile 2-ml tubes containing 500 μ l of guanidine isothiocyanate (GITC) buffer [5 mol l⁻¹GITC, 100 mmol l⁻¹EDTA (pH 8.0), 0.5% Sarkosyl] and stored overnight at -80 °C. DNA was extracted using DNeasy blood and tissue kit (Qiagen, Inc., Valencia, CA). Extracted DNA were resuspended in 200 μ l AE buffer, and stored at -80 °C until processed.

For the HS-esp marker, 125 ml of each sewage spiked water sample was filtered through 0.45- μ m pore size membranes (Advantec). The membranes were then transferred to mEI agar (Difco, Detroit, MI) and incubated at 41 °C for 48 h. After incubation, the filter papers were suspended in tryptic soy broth (TSB) (Oxoid, London, UK) and incubated at 41 °C for 3 h for enrichment (Scott et al., 2005). DNA was extracted from 2 ml of enriched culture by using QIAamp stool DNA kit (Qiagen) according to the manufacturer's instructions. Extracted DNA was resuspended in 200 μ l AE buffer, and stored at –80 °C until processed.

For viral markers (HS-AVs and HS-PVs) and enteric viruses (i.e., EVs, SVs and TVs), each dilution was concentrated by using the previously published method (Haramoto et al., 2005). Briefly 125 ml of each sewage spiked water sample was supplemented with 2.5 mM MgCl₂ and then passed through an HA electronegative filter (0.45- μ m pore size, 90 mm diameter; Millipore, Tokyo, Japan) attached to a glass filter holder

TargetsPrimer sequence (5'-3')Amplice size (bs)Number of SCR (cvcs)ReferenceHuman-specific BacteroidesF - ATC ATG AGT TCA CAT GTC GG52045Bernhard and Field, 2000Cattle-specific BacteroidesF - CCA ACY TC CCG WTA CTC58045Bernhard and Field, 2000CF128 (CS-CF12)R - CAA TCG GAG TTC TC GT680455Scott et al., 2005Human-specific enterococciF - TAT GAA AGC AAC GAC AGT ACT CC680450Scott et al., 2005Human-specific enterococciF - CAC GCG ATG GTC TTA CAT GCA CAT CC300900Forg et al., 2005Human-specific polyR - CAC GCA GTG GTC TTA CAT GCA CAT CC300900Forg et al., 2005Human-specific polyR - CAC GCA GCG GTA GTC TA CAT GCA CAT CC143Human-specific polyR - GAG CAT GCG GAG TA CTA CAG1721000McQuaig et al., 2005Human-specific polyR - GAG CCT AGC GAA GCT AGC GCA GCA1721000Haramote et al., 2005Human-specific polyF - CGG GTG CCG DAG GTA GTA CAG7945Haramote et al., 2005Human-specific polyF - CGG GCT GCG CAAT CCAAC19650Haramote et al., 2005Human-specific polyF - CGA GG CT GC CCA CT AGC19650Haramote et al., 2005Human-specific polyF - CGA GG CT GC CAC CT AGC19650Haramote et al., 2005Human-specific polyF - CGA GG CT GC CCA CT AGC19650Haramote et al., 2005R - GCGF - CCC GG CT CT CG CAC CT AGC19650	Table 1 – Primer sequence used for the detection of host-specific molecular markers and enteric viruses.							
Human-specific Bacteroides HF183 (HS-HF183)F - ATC ATG AGT TCA CAT GTC CG52045Bernhard and Field, 2000Cattle-specific BacteroidesF - CAA ATCG GAG TTC TTC GTG58045Bernhard and Field, 2000CF128 (CS-CF12)R - CAA TCG GAG TTC TTC GTG68045Scott et al., 2005Human-specific enterococciF - TAT GAA AGC AAC AGC ACA AGT T CGA68045Scott et al., 2005surface protein (esp) marker (HS-esp)R - ACG TCG AAA GTT CGA TTT CC30090Fong et al., 2005Human-specific adenoviruses (HS-AVs)F - GCC GCA GTG GTC TTA CAT GCA CAT C R - CAG CAC GCG GAG TAC TTC AGC CTG143-Human-specific polyomaviruses (HS-PVS)F - GGT GCC AAC CTA TGG AAC AG172100McQuaig et al., 2006HS-PVS)R - GGT GCC AAC CTA TGG ACA TG R - GGT GCC GCG CGG GA GTG ACA AG7945Haramoto et al., 2005F - GCG CCC CGG GCG CCT GAA TG (HS-PVS)F - CGC GCC CCT GAA TG R - GGT GCC AATC CAA7941Haramoto et al., 2005Enteroviruses (IVs)F - CCT CCG GCC CT GAA TG R - ACC GGA TGG CCA ATC CAA19650Haramoto et al., 2008Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC F3 - TTT GGC CCT CCA TTC CAAA CAC TAT TTTG F2 - CCA GGC TCT CCG CAC CTAC F3 - TTT GGC CCT CCA CTAC F3 - TTT GGC CCT CAC TTC CAAA CAC TAA TTTG19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC F3 - TTT GGC CCT CCA CTAC F3 - TTT GGC CCT CCA CTAC F3 - CCA GGC CTC CAC CTAC F3 - TTT GGC CCT CC	Targets	Primer sequence (5'- 3')	Amplicon size (bp)	Number of PCR cycles	Reference			
HF183 (HS-HF183)R - CAA TCG GAG TTC TTC GTGCattle-specific BacteroidesF - CCA ACY TTC CCG WTA CTC58045Bernhard and Field, 2000CF128 (CS-CF12)R - CAA TCG GAG TTC TTC GTG58045Scott et al., 2005Human-specific enterococciF - TAT GAA AGC AAC AGC ACA AGT T68045Scott et al., 2005surface protein (esp) marker (HS-esp)R - ACG TCG AAA GTT CGA TTT CC30090Fong et al., 2005Human-specific adenoviruses (HS-AVs)R - GCC GCA GTG GTC TTA CAT GCA CAT C30090Fong et al., 2005(HS-AVs)R - CAG CAC GCC GCG GAT GTC AAA GT F - GCC ACC GAG ACG TAC CTC AGC CTG143Human-specific polyomaviruses (HS-PVs)F - AGT CTT TAG GGT CTT CTA CC172100McQuaig et al., 2006(HS-PVs)R - GGT GCC AAC CTA TGG AAC AGTTorquetenoviruses (TVs)F - CCT CCG GCC CCT GAA TG19650Haramoto et al., 2005R - ACC GGA TGG CCT CGC CAC CTAC CAAF1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC CTC CGT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTACF3 - TT GGC CCT CGT CAT CTC CAA CAC TAT TTTGF2 - CCC GGC CTC CAT CTC AAA CAC TAT TTTGF3 - TTT GGC CTT CAA CAC TACF3 - TTT GGC CTT CAA CAC TAC	Human-specific Bacteroides	F – ATC ATG AGT TCA CAT GTC CG	520	45	Bernhard and Field, 2000			
Cattle-specific BacteroidesF - CCA ACY TTC CCG WTA CTC58045Bernhard and Field, 2000CF128 (CS-CF12)R - CAA TCG GAG TTC TTC GTG845Scott et al., 2005Human-specific enterococciF - TAT GAA AGC AAC AGC AGC AGT TC CC68045Scott et al., 2005surface protein (esp) markerR - ACG TCG AAA GTT CGA TT CC30090Fong et al., 2005(HS-esp)F - GCC GCA GTG GTC TTA CAT GCA CAT C30090Fong et al., 2005Human-specific adenovirusesF - GCC GCG GAT GTC AAA GT143143R - TTG TAC GAG TAC GCG GTA CTC AGC CTG143144144R - TTG TAC GAG TAC CCG GTA TCT ACC CCG CGG TC172100McQuaig et al., 2006Human-specific polyomavirusesF - AGT CTT AGG GTG CTT ACAC CAG7945Haramoto et al., 2005R - GGT GCC AAC CTA TGG GAC ATG7945Haramoto et al., 2005143Cruptetenoviruses (TVs)F - CGG GTG CCG DAG GTG AGT TTA CAC7945Haramoto et al., 2005R - ACC GGA TGG CCA TRG CCC GGC CAG19650Haramoto et al., 2005R - ACC GGA TGG CCA TC CGC CAC CTAC19650Haramoto et al., 2008Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC CTT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTACF4 - GCC GC CTC CAT TTC AAA CAC TAT TTTGF5 - GCC GC CTC CAT TTC AAA CAC TAC <td>HF183 (HS-HF183)</td> <td>R – CAA TCG GAG TTC TTC GTG</td> <td></td> <td></td> <td></td>	HF183 (HS-HF183)	R – CAA TCG GAG TTC TTC GTG						
CF128 (CS-CF12)R - CAA TCG GAG TTC TTC GTGHuman-specific enterococciF - TAT GAA AGC AAC AGC ACA AGT T68045Scott et al., 2005surface protein (esp) marker (HS-esp)R - ACG TCG AAA GTT CGA TTT CC30090Fong et al., 2005Human-specific adenovirusesF - GCC GCA GTG GTC TTA CAT GCA CAT C30090Fong et al., 2005(HS-AVs)R - CAG CAC GCC GCG GAT GTC AAA GT143R - TTG TAC GAG TAC GCG GTA TCC TCG CGG TC143R - TTG TAC GAG TAC GCG GTA TCC TCG CGG TC172100McQuaig et al., 2006(HS-PVs)R - GGT GCC AAC CTA TGG AAC AG7945Haramoto et al., 2005R - GGG CCT CCG CCG CCG GGA TGC CAAC7945Haramoto et al., 2005R - GAG CCT TGC CCA TTG CCA ATG19650Haramoto et al., 2005R - ACG GAG CTC CGC CAC CTAC19650Haramoto et al., 2008R - CCG GCT CCG CAC CTAC CTAC19650Haramoto et al., 2008R - CCG GCT CCG CAC CTAC CTAC19650Haramoto et al., 2008R - ACC GGC TCT CGC CAC CTAC19650Haramoto et al., 2008R - ACC GGC TCT CGC CAC CTAC19650Haramoto et al., 2008R - CCA GCC TC CGC CAC CTAC19650Haramoto et al., 2008R - GCC CTC CAT CTC CAA CCTAC19650Haramoto et al., 2008R - GCC CTC CAT CTC CAA CAC CTAC19650Haramoto et al., 2008R - GCC CTC CAT CTC CAA CAC CTAC19650Haramoto et al., 2008R - GCC CTC CAT CTC CAA CAC CTAC196 </td <td>Cattle-specific Bacteroides</td> <td>F – CCA ACY TTC CCG WTA CTC</td> <td>580</td> <td>45</td> <td>Bernhard and Field, 2000</td>	Cattle-specific Bacteroides	F – CCA ACY TTC CCG WTA CTC	580	45	Bernhard and Field, 2000			
Human-specific enterococciF - TAT GAA AGC AAC AGC AGC AGA AGT T68045Scott et al., 2005surface protein (esp) marker (HS-esp)R - ACG TCG AAA GTT CGA TTT CC30090Fong et al., 2005Human-specific adenoviruses (HS-AVs)F - GCC GCA GTG GTC TTA CAT GCA CAT C30090Fong et al., 2005Muman-specific polyomaviruses (HS-AVs)F - GCC GCA GTG GTC TTA CAT GCA CAT C143R - TTG TAC GAG TAC GCG GTA TCC TCG CGG TC143R - TTG TAC GAG TAC GCG GTA TCC TCG CGG TC172100McQuaig et al., 2006(HS-PVs)R - GGT GCC AAC CTA TGG AAC AG7945Haramoto et al., 2005Corquetenoviruses (TVs)F - CCG GTG CCG DAG GTG AGT TTA CAC7945Haramoto et al., 2005Enteroviruses (EVs)F - CCT CCG GCC CCT GAA TG19650Haramoto et al., 2008Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTAC	CF128 (CS-CF12)	R – CAA TCG GAG TTC TTC GTG						
surface protein (esp) marker (HS-esp)R - ACG TCG AAA GTT CGA TTT CCHuman-specific adenoviruses (HS-AVs)F - GCC GCA GTG GTC TTA CAT GCA CAT C30090Fong et al., 2005(HS-AVs)R - CAG CAC GCC GCG GAT GTC AAA GT F - GCC ACC GAG ACG TAC TTC AGC CTG143143(HS-PVs)R - TTG TAC GAG TAC GCG GTA TCC TCG CGG TC172100McQuaig et al., 2006(HS-PVs)R - GGT GCC AAC CTA TGG AAC AG7945Haramoto et al., 2005(HS-PVs)F - CCT CCG GCC CCT GAA TG19650Haramoto et al., 2005R - GAG CCT TGC CCA TTC GC CAC CTAC19650Haramoto et al., 2008F - CCT CCG GCC CTC GC CAC CTAC CTAC19650Haramoto et al., 2008Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC196	Human-specific enterococci	F – TAT GAA AGC AAC AGC ACA AGT T	680	45	Scott et al., 2005			
Human-specific adenoviruses (HS-AVs)F - GCC GCA GTG GTC TTA CAT GCA CAT C30090Fong et al., 2005(HS-AVs)R - CAG CAC GCC GCG GAT GTC AAA GT F - GCC ACC GAG ACG TAC TTC AGC CTG143R - TTG TAC GAG TAC GCG GTA TCC TCG CGG TC143Human-specific polyomavirusesF - AGT CTT TAG GGT CTT CTA CC172100McQuaig et al., 2006 (HS-PVs)R - GGT GCC AAC CTA TGG AAC AG7945Torquetenoviruses (TVs)F - CGG GTG CCG DAG GTG AGT TTA CAC7945Haramoto et al., 2005R - GAG CCT TGC CCA TRG CCC GGC CAG19650Enteroviruses (EVs)F - CCT CCG GCC CAC CTAC19650R - ACC GGA TGG CCA ATC CAA19650Haramoto et al., 2008Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650F2 - CCA GGC TCT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTAC19650F3 - TTT GGC CCT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTAC19650F4 - GCC CTC CAT TTC AAA CAC TAT TTTGF2 - GCC CTC CAT TTC CAAA CAC TAT TTTG100	surface protein (esp) marker (HS-esp)	R – ACG TCG AAA GTT CGA TTT CC						
(HS-AVs)R - CAG CAC GCC GCG GAT GTC AAA GTF - GCC ACC GAG ACG TAC TTC AGC CTG143R - TTG TAC GAG TAC GCG GTA TCC TCG CGG TCHuman-specific polyomavirusesF - AGT CTT TAG GGT CTT CTA CC(HS-PVs)R - GGT GCC AAC CTA TGG AAC AGTorquetenoviruses (TVs)F - CGG GTG CCG DAG GTG AGT TTA CACP - GAG CCT TGC CCA TRG CCC GGC CAG7945Haramoto et al., 2005R - GAG CCT TGC CCA TRG CCC GGC CAGEnteroviruses (EVs)F - CCT CCG GCC CT GAA TG9509Haramoto et al., 20058- ACC GGA TGG CCA ATC CAASapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC9509Haramoto et al., 20089509Haramoto et al., 200895091495091495091495091495091495091495091495091491091091091091091091091091091091091091091091091091010 <t< td=""><td>Human-specific adenoviruses</td><td>F – GCC GCA GTG GTC TTA CAT GCA CAT C</td><td colspan="2">300 90</td><td colspan="2">Fong et al., 2005</td></t<>	Human-specific adenoviruses	F – GCC GCA GTG GTC TTA CAT GCA CAT C	300 90		Fong et al., 2005			
F - GCC ACC GAG ACG TAC TTC AGC CTG143R - TTG TAC GAG TAC GCG GTA TCC TCG CGG TCHuman-specific polyomavirusesF - AGT CTT TAG GGT CTT CTA CC172100McQuaig et al., 2006(HS-PVs)R - GGT GCC AAC CTA TGG AAC AG7945Haramoto et al., 2005Torquetenoviruses (TVs)F - CGG GTG CCG DAG GTG AGT TTA CAC7945Haramoto et al., 2005R - GAG CCT TGC CCA TRG CCC GGC CAG19650Haramoto et al., 2005Enteroviruses (EVs)F - CCT CCG GCC CT GAA TG19650Haramoto et al., 2008Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC CTC CAT TTC AAA CAC TAT TTTG143143F3 - TTT GGC CCT CGC CAC CTAC143143F3 - TTT GGC CCT CAT TTC AAA CAC TAA TTTG145143	(HS-AVs)	R – CAG CAC GCC GCG GAT GTC AAA GT						
R - TTG TAC GAG TAC GCG GTA TCC TCG CGG TCHuman-specific polyomavirusesF - AGT CTT TAG GGT CTT CTA CC172100McQuaig et al., 2006(HS-PVs)R - GGT GCC AAC CTA TGG AAC AG7945Haramoto et al., 2005Torquetenoviruses (TVs)F - CGG GTG CCG DAG GTG AGT TTA CAC7945Haramoto et al., 2005R - GAG CCT TGC CCA TRG CCC GGC CAG19650Haramoto et al., 2005Enteroviruses (EVs)F - CCT CCG GCC CT GAA TG19650Haramoto et al., 2005Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F3 - TTT GGC CCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F3 - TTT GGC CCT CGC CAC CTAC19650Haramoto et al., 2008F3 - TTG GC CCT CGC CAC CTAC19650Haramoto et al., 2008F3 - TTG GC CCT CGC CAC CTAC19650Haramoto et al., 2008F3 - TTG GC CCT CAT TTC AAA CAC TAT TTTG19650Haramoto et al., 2008F4 - GCC CTC CAT TTC AAA CAC TAA TTTG19650Haramoto et al., 2008F5 -		F – GCC ACC GAG ACG TAC TTC AGC CTG	143					
Human-specific polyomavirusesF - AGT CTT TAG GGT CTT CTA CC172100McQuaig et al., 2006(HS-PVs)R - GGT GCC AAC CTA TGG AAC AG7945Haramoto et al., 2005Torquetenoviruses (TVs)F - CGG GTG CCG DAG GTG AGT TTA CAC7945Haramoto et al., 2005R - GAG CCT TGC CCA TRG CCC GGC CAG19650Haramoto et al., 2005Enteroviruses (EVs)F - CCT CCG GCC CT GAA TG19650Haramoto et al., 2005Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - GCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - GCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - GCC CTC CAT TTC AAA CAC TAT TTTG777777F3 - TTT GGC CCT CAT TTC AAA CAC TAA TTTG777777F3 - TTC GGC CTC CAT TTC AAA CAC TAA TTTG777777		R – TTG TAC GAG TAC GCG GTA TCC TCG CGG TC						
(HS-PVs)R - GGT GCC AAC CTA TGG AAC AGTorquetenoviruses (TVs)F - CGG GTG CCG DAG GTG AGT TTA CAC7945Haramoto et al., 2005R - GAG CCT TGC CCA TRG CCC GGC CAG19650Haramoto et al., 2005Enteroviruses (EVs)F - CCT CCG GCC CCT GAA TG19650Haramoto et al., 2005R - ACC GGA TGG CCA ATC CAA19650Haramoto et al., 2008Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - GCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - GCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - GCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - GCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F3 - TTT GGC CCT CGC CAC CTAC19650Haramoto et al., 2008F2 - GCC CTC CAT TTC AAA CAC TAT TTTG19650F3 - TTT GGC CCT CAT TTC AAA CAC TAA TTTG196106	Human-specific polyomaviruses	ific polyomaviruses F – AGT CTT TAG GGT CTT CTA CC		100	McQuaig et al., 2006			
Torquetenoviruses (TVs)F - CGG GTG CCG DAG GTG AGT TTA CAC7945Haramoto et al., 2005R - GAG CCT TGC CCA TRG CCC GGC CAGF - CCT CCG GCC CCT GAA TG19650Haramoto et al., 2005Enteroviruses (EVs)F - CCT CCG GCC CCT GAA TG19650Haramoto et al., 2005Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTACR1 - GCC CTC CAT CTC AAA CAC TAT TTTGR2 - GCC CTC CAT TTC AAA CAC TAA TTTG	(HS-PVs)	R – GGT GCC AAC CTA TGG AAC AG						
R - GAG CCT TGC CCA TRG CCC GGC CAG Enteroviruses (EVs) F - CCT CCG GCC CT GAA TG 196 50 Haramoto et al., 2005 R - ACC GGA TGG CCA ATC CAA Sapoviruses (SVs) F1 - CCA GGC TCT CGC CAC CTAC 196 50 Haramoto et al., 2008 F2 - CCA GGC TCT CGC TAC CTAC F3 - TTT GGC CCT CGC CAC CTAC R1 - GCC CTC CAT CTC AAA CAC TAT TTTG R2 - GCC CTC CAT TTC AAA CAC TAA TTTG	Torquetenoviruses (TVs)	F – CGG GTG CCG DAG GTG AGT TTA CAC	79	45	Haramoto et al., 2005			
Enteroviruses (EVs)F - CCT CCG GCC CCT GAA TG19650Haramoto et al., 2005R - ACC GGA TGG CCA ATC CAAR - ACC GGA TGG CCA ATC CAA19650Haramoto et al., 2008Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTACR1 - GCC CTC CAT CTC AAA CAC TAT TTTGR2 - GCC CTC CAT TTC AAA CAC TAA TTTGF3 - TTT GGC CTC CAT TTC AAA CAC TAA TTTGF3 - TTT GGC CTC CAT TTC AAA CAC TAA TTTG		R – GAG CCT TGC CCA TRG CCC GGC CAG						
R - ACC GGA TGG CCA ATC CAA Sapoviruses (SVs) F1 - CCA GGC TCT CGC CAC CTAC 196 50 Haramoto et al., 2008 F2 - CCA GGC TCT CGC TAC CTAC F3 - TTT GGC CCT CGC CAC CTAC R1 - GCC CTC CAT CTC AAA CAC TAT TTTG R2 - GCC CTC CAT TTC AAA CAC TAA TTTG	Enteroviruses (EVs)	F – CCT CCG GCC CCT GAA TG	196 50		Haramoto et al., 2005			
Sapoviruses (SVs)F1 - CCA GGC TCT CGC CAC CTAC19650Haramoto et al., 2008F2 - CCA GGC TCT CGC TAC CTACF3 - TTT GGC CCT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTACR1 - GCC CTC CAT CTC AAA CAC TAT TTTGR2 - GCC CTC CAT TTC AAA CAC TAA TTTGF3 - TTT GGC CCT CGC CAC CTACF3 - TTT GGC CCT CGC CAC CTAC		R – ACC GGA TGG CCA ATC CAA						
F2 – CCA GGC TCT CGC TAC CTAC F3 – TTT GGC CCT CGC CAC CTAC R1 – GCC CTC CAT CTC AAA CAC TAT TTTG R2 – GCC CTC CAT TTC AAA CAC TAA TTTG	Sapoviruses (SVs)	F1 – CCA GGC TCT CGC CAC CTAC	196 50		Haramoto et al., 2008			
F3 – TTT GGC CCT CGC CAC CTAC R1 – GCC CTC CAT CTC AAA CAC TAT TTTG R2 – GCC CTC CAT TTC AAA CAC TAA TTTG		F2 – CCA GGC TCT CGC TAC CTAC						
R1 – GCC CTC CAT CTC AAA CAC TAT TTTG R2 – GCC CTC CAT TTC AAA CAC TAA TTTG		F3 – TTT GGC CCT CGC CAC CTAC						
R2 – GCC CTC CAT TTC AAA CAC TAA TTTG		R1 – GCC CTC CAT CTC AAA CAC TAT TTTG						
		R2 – GCC CTC CAT TTC AAA CAC TAA TTTG						

(Advantec, Tokyo, Japan). Subsequently, 100 ml of 0.5 mM H_2SO_4 (pH 3) was passed through the filter to remove magnesium ions and other electropositive substances, followed by filtration of 10-ml of 1 mM NaOH (pH 11) for elution of viruses from the filter. The filtrate was recovered in a tube containing 100- μ l of 100 mM H₂SO₄ (pH 1) and 100- μ l of 100× Tris-EDTA buffer (pH 8) for neutralization. All 10-ml elutes were stored at -20 °C until further processing. The concentrated samples were further purified, concentrated, and desalted with Centriprep YM-50 concentrator columns (Millipore). Samples were added to the Centriprep YM-50 and centrifuged at 1000g for 10 min, followed by removal of the sample that passed through the ultrafiltration membrane (10 ml) and further centrifugation at 1000 g for 10 min to obtain a final volume of 700-µl. Concentrates were split in half (2 \times 350 μ l), and stored at -80 °C. Viral DNA and RNA were extracted from each concentrated samples using DNeasy blood and tissue kit and QIAamp viral RNA mini kit (Qiagen) according to the manufacture's instructions. For viral DNA (HS-AVs, HS-PVs, TVs) and RNA (EVs and SVs) extractions respectively, 200 μ l and 140 μ l of concentrated samples were used. Extracted viral DNA was resuspended in 200 µl AE buffer, and extracted viral RNA was resuspended in 60 µl AVE buffer, respectively, and stored at -80 °C until processed.

2.7. Real-time PCR positive controls

The HS-HF183, HS-AVs, HS-PVs, EVs, SVs and TVs positive controls were isolated from sewage. In brief, the real-time PCR-amplified product was purified using a QIAquick PCR purification kit (Qiagen) and cloned into a pGEM-T Easy vector system (Promega, Madison, WI), transferred into *E. coli* JM109competent cells, and plated on LB agar plates containing ampicillin, isopropyl-β-D-thiogalactopyranoside, and 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside, as recommended by the manufacturer. Recombinant plasmids with corresponding inserts were purified using a plasmid mini kit (Qiagen). DNA sequencing was carried out at the Australian Genome Research Facility (St. Lucia, Queensland, Australia). For the HS-esp marker, genomic DNA was extracted from a positive control (*E. faecium* C68 strain, provided by Dr. Louis B. Rice of the Louis Stokes Cleveland veterans Affairs Medical Centre in Cleveland, OH, USA).

2.8. Real-time PCR analysis for bacteria, DNA and RNA viruses

The real-time PCRs were performed using a Rotor-Gene 6000 real-time cycler (Corbett Research, Mortlake, Australia). Amplification was performed in 25- μ l reaction mixtures using Platinum SYBR Green qPCR SuperMix-UDG (Invitrogen, Carlsbad, CA). The HS-HF183 and the HS-*esp* real-time PCR mixture contained 12.5 μ l of SuperMix, 300 nM of each primer, and 5 μ l of DNA extract. For both HS-HF183 and the HS-*esp* markers, the real-time PCR consisted of 2 min at 50 °C, 15 min at 95 °C followed by 45 cycles of 30 s at 94 °C, 60 s at 59 °C and 60 s at 72 °C.

For the HS-AVs detection a nested real-time PCR protocol was used (Fong and Lipp, 2005). The first round of real-time PCR contained 12.5 μ l of SuperMix, 300 nM of each primer, and 5 μ l of DNA extract. The first round of real-time PCR consisted of 2 min at 50 °C, 4 min at 94 °C followed by 45 cycles of 30 s at 92 °C, 30 s at 60 °C and 60 s at 72 °C, and a final extension of 5 min at 72 °C. The second round of real-time PCR was run under the same reaction conditions and cycling parameters described above, with 3 μ l of product from the first round as the template.

A nested real-time PCR protocol was also used for HS-PVs detection. The first round of real-time PCR contained 12.5 μ l of SuperMix, 200 nM of each primer, and 5 μ l of DNA extract. The first round of real-time PCR consisted of 2 min at 50 °C,

2 min at 94 °C followed by 50 cycles of 20 s at 94 °C, 20 s at 55 °C and 20 s at 72 °C, and a final extension of 2 min at 72 °C. The second round of real-time PCR was run under the same reaction conditions and cycling parameters described above, with 3 μ l of product from the first round as the template. TVs real-time PCR contained 12.5 μ l of SuperMix, 400 nM of each primer, and 5 μ l of template DNA. The real-time PCR consisted of 2 min at 50 °C, 10 min at 95 °C followed by 45 cycles of 10 s at 95 °C, 30 s at 62 °C.

For the detection of EVs and SVs, 8 μ l of extracted RNA was added to a reaction mixture containing 1 μ l of annealing buffer and 1 μ l of 50 ng/ μ l random hexamers and was incubated at 65 °C for 5 min, then quenched at 4 °C for 60 s. Subsequently, 10 μ l of 2× reaction mix, 2 μ l SuperScript III enzyme mix (Invitrogen), then were heated at 25 °C for 10 min then 50 °C for 50 min. The reaction was terminated with a 85 °C incubation for 5 min. EVs real-time PCR contained 12.5 μ l of Super-Mix, 400 nM of each primer, and 2 μ l of cDNA. The real-time PCR consisted of 2 min at 50 °C, 10 min at 95 °C followed by 50 cycles of for 15 s at 95 °C, 60 s at 60 °C. SVs real-time PCR contained 12.5 μ l of SuperMix, 400 nM of each primer, and 5 μ l of cDNA. The real-time PCR consisted of 2 min at 50 °C, 10 min at 95 °C followed by 50 cycles of 15 s at 95 °C, 60 s at 56 °C.

2.9. Real-time PCR quality control

For each real-time PCR experiment, corresponding positive (i.e., target DNA) and negative (sterile water) controls were included. Each DNA sample was tested in triplicate to obtain positive/negative results. To separate the specific product from non-specific products (if any), DNA melting curve analysis was performed for each real-time PCR experiment. During melting curve analysis, the temperature was increased from 53 to 95 °C at approximately 2 °C/min. Samples were considered positive when the visible band was the same as that of the positive control strain and had the same melting temperature as the positive control. To minimise real-time PCR set up were performed in separate laboratories. To prevent false positive results for water samples, a method blank was included.

2.10. Real-time PCR detection limit

To determine the real-time PCR detection limits, genomic DNA and plasmid DNA were quantified using a spectrophotometer. Ten-fold serial dilutions (i.e., 10^3-10^0) were made using buffer AE (Qiagen) for each target and tested with the real-time PCR. The lowest concentration of gene copies detected consistently in replicate assays was considered as the real-time PCR detection limit.

3. Results

3.1. Specificity of the real-time PCR primers

Primer specificity was determined by searching for similar sequences in microbial genomes using the Basic Local Alignment Search Tool (BLAST) program (http://www.ncbi.nlm.nih. gov/BLAST/). No homology was observed with known gene sequences of other pathogenic microorganisms commonly found in environmental waters.

3.2. Real-time PCR inhibitors

To detect the presence of inhibitors, DNA of animal fecal and STP samples was spiked with 10^3 gene copies the HS-HF183 and CS-CF128 markers. The real-time PCR C_T values were compared to those obtained from the same concentrations of DNA that was used to spike 1 L of distilled water. For the spiked distilled water, the mean C_T values for the HS-HF183, HS-*esp*, HS-AVs and HS-PVs are shown in Table 2.

For STP DNA samples (extracted for the HS-HF183 specificity and sensitivity assay), the mean $C_{\rm T}$ values were 28 ± 0.5 when undiluted DNA was spiked with cattle-specific CF128 markers. For 10-fold and 100-fold dilutions of DNA, these values were 23 ± 0.7 , and 23 ± 0.5 , respectively. Similarly, for animal fecal DNA samples (extracted for the HS-HF183 specificity and sensitivity assay), the mean $C_{\rm T}$ values were 26 ± 1.1 when undiluted DNA was spiked with the HS-HF183 markers. For 10-fold and 100-fold dilutions of DNA, these values were

Table 2 – Effects of PCR inhibitors on the real-time PCR
detection of spiked human-specific HS-HF183 and
CS-CF128 markers in animal fecal and sewage samples
as opposed to distilled water samples.

Undiluted DNA10-fold dilution100-fold dilutionHS-HF183 host-specificity assay $ -$ Distilled water 23 ± 0.3 $ -$ STP (primary influent) 28 ± 0.5 23 ± 0.7 23 ± 0.5 Distilled water 21 ± 0.3 $ -$ Cottle 21 ± 0.3 $ -$								
HS-HF183 host-specificity assay Distilled water 23 ± 0.3 STP (primary 28 ± 0.5 23 ± 0.7 23 ± 0.5 influent) Distilled water 21 ± 0.3 Cottle 25 ± 0.1 21 ± 1.2 21 ± 0.7								
Distilled water 23 ± 0.3 STP (primary 28 ± 0.5 23 ± 0.7 23 ± 0.5 influent) Distilled water 21 ± 0.3 Cottle 25 ± 0.1 21 ± 1.2 21 ± 0.7								
STP (primary influent) 28 ± 0.5 23 ± 0.7 23 ± 0.5 Distilled water 21 ± 0.3 Cottle 25 ± 0.1 21 ± 1.2 21 ± 0.7								
Distilled water 21 ± 0.3								
Cattle 25 ± 0.1 21 ± 1.2 21 ± 0.7								
Pig 27 ± 0.7 22 ± 0.1 23 ± 1.2								
$Sheep \qquad 26 \pm 0.1 \qquad 22 \pm 0.4 \qquad 22 \pm 0.9$								
$Dog \qquad 24 \pm 2.0 \qquad 22 \pm 0.3 \qquad 22 \pm 0.2$								
Duck 26 ± 0.4 21 ± 7.0 21 ± 1.0								
HS-esp host-specificity assay								
Distilled water 23 ± 0.7 – –								
STP (primary 24 ± 0.2 23 ± 0.9 – influent)								
Distilled water 22 ± 0.6 – –								
Cattle 23 ± 0.4 22 ± 0.9 –								
Pig 23 ± 0.9 23 ± 0.9 –								
Sheep 22 ± 1.2 22 ± 0.8 –								
Dog 23 ± 0.6 23 ± 0.9 –								
Duck 24 ± 0.2 23 ± 0.7 –								
HS-AVs and HS-PVs host-specificity assays								
Distilled water 26 ± 0.3 – –								
STP (primary 26 ± 0.9 26 ± 0.6 – influent)								
Distilled water 25 ± 0.4 – –								
Cattle 26 ± 0.7 26 ± 0.4 –								
Pigs 26 ± 0.3 25 ± 0.8 –								
Sheep 25 ± 0.9 25 ± 0.3 –								
Dogs 27 ± 0.3 26 ± 0.7 –								
Ducks 26 ± 0.6 25 ± 0.2 –								

 22 ± 0.5 , and 22 ± 0.8 , respectively. One-way ANOVA was performed to determine the differences between the C_T values obtained for distilled water and those obtained for animal fecal DNA samples extracted for the HS-HF183 assay. Significant (P < 0.001) differences were observed between the C_T values for spiked distilled water and undiluted DNA from animal fecal samples, indicating that the undiluted DNA extracted from animal fecal samples contained PCR inhibitory substances. However, no significant differences (P > 0.05) were observed between the C_T values for spiked distilled water and serially diluted DNA (i.e., 10-fold and 100-fold) indicating that 10-fold dilution of DNA is required to remove the effects of real-time PCR inhibitory substances from animal fecal samples. For STP and animal fecal DNA samples (extracted for the HS-esp, HS-AVs and HS-PVs specificity and sensitivity assays), the mean C_T values for undiluted DNA were similar to the values obtained for distilled water indicating the samples were inhibitor free.

3.3. Host-specificity and sensitivity of human-specific molecular markers

Host-specificity is the probability to detect when a source is not present and sensitivity is the probability to detect a source when it is present. The HS-HF183 marker was detected in all sewage samples and was not detected in 50 animal fecal samples (Table 3). Only one fecal sample from a dog was found to be positive. The overall host-specificity of the HS-HF183 marker to differentiate between humans and animals was 98%, and the overall sensitivity of these markers in sewage samples was 100%. Similarly, the HS-esp marker was detected in all STP samples, and was not detected in any animal fecal sample. The overall host-specificity and sensitivity of the HSesp markers were 100%. The HS-AVs were detected in 31 out of 32 STP samples, and HS-PVs were detected in all STP samples. However, these markers could not be detected in any of the 50 animal fecal samples tested. The overall specificity of both HS-AVs and PVs was 100%. The overall sensitivity of the HS-AVs and HS-PVs was 97% and 100%, respectively.

Table 3 – Polymerase chain reaction positive results for human-specific Bacteroides HF183 (HS-HF183), humanspecific Enterococci faecium enterococci surface protein markers (HS-esp), human-specific adenoviruses (HS-AVs) and human-specific polyomaviruses (HS-PVs) in host groups in Southeast Queensland, Australia.

Host groups	No. of samples tested/real-time positive results			
	HS-HF183	HS-esp	HS-AVs	HS-PVs
STP (primary influent)	32/32	32/32	32/31	32/32
Cattle	10/0	10/0	10/0	10/0 ^a
Pigs	10/0	10/0	10/0	10/0 ^a
Sheep	10/0	10/0	10/0	10/0 ^a
Dogs	10/1	10/0	10/0	10/0
Ducks	10/0	10/0	10/0	10/0

a Composite wastewater samples.

3.4. Real-time PCR limit of detection

The real-time PCR limits of detection were performed by analysing purified plasmid/genomic DNA isolated from the bacterial and viral targets. To determine the reproducibility of the assay, several replicates (n = 10) were tested. The results of these assays are summarised in Table 4. The realtime PCR detection limits were as low as one gene copy per reaction for the HS-HF183 and HS-*esp* markers. For HS-AVs, HS-PVs, EVs, SVs and TVs, the detection limits were ten gene copies per reaction. Lower levels (i.e., one copy) were detected, but the results were not reproducible for all replicates.

3.5. Concentrations of fecal indicators in sewage spiked water samples

The concentration of E. coli in sewage spiked freshwater dilution of 10^{-1} (i.e., contains 6.25 ml of sewage) was $4.3 \pm 0.3 \times 10^5$ colony forming units (CFUs) (Table 5). For enterococci and C. *perfringens* spores these figures were $9.6 \pm 0.2 \times 10^4$ and $5.0 \pm 1.0 \times 10^2$, respectively. Culturable E. coli were detected at dilution up to 10^{-6} and the concentration of E. coli at this dilution was $3.0 \pm 2.0 \times 10^0$ CFU. Enterococci and C. *perfringens* spores were detected at dilutions of up to 10^{-5} and 10^{-3} respectively, and the concentrations at these dilutions were $7.1 \pm 3.0 \times 10^0$ and $5.1 \pm 3.0 \times 10^0$. The concentration of E. coli was higher than enterococci and C. *perfringens* spores.

3.6. The real-time PCR ability to detect fresh sewage in sewage spiked water samples

For freshwater, the real-time assay resulted in the detection of the HS-HF183 marker up to dilution 10^{-8} . At this dilution, no culturable E. coli, enterococci and C. perfringens spores were found. The HS-esp marker was detected up to dilution 10^{-4} and the concentration of E. coli and enterococci at this dilution were 3.9 ± 0.8 \times 10^2 and 8.1 ± 6.0 \times 10^1 CFUs. However, no *C. perfringens* spores were found at this dilution. The HS-AVs and HS-PVs were detected up to dilution 10⁻⁴ by nested real-time PCR protocol. Real-time PCR assays of enteric viruses were similar to HS-esp, HS-AVs and HS-PVs. However, EVs were detected up to dilution 10^{-5} and the concentrations of E. coli and enterococci at this dilution were only $4.0\pm1.0\,\times\,10^1$ and $7.1\pm6.0\,\times\,10^1$ CFUs. The ability of real-time PCR assays of human-specific markers and enteric viruses for seawater and distilled water was similar to that of freshwater (see Table 5).

4. Discussion

The host-specificity and sensitivity of the human-specific markers were assessed by testing fecal DNA samples from target (i.e., sewage samples) and non-target (i.e., animal fecal samples) sources. The HS-HF183 marker was detected in one animal fecal sample out of 50 samples tested. Previous studies also report the presence of the host-specific *Bacteroides* markers in a small number of non-target samples (Ahmed

Table 4 – Real-time PCR limit of detection for host-specific molecular markers and enteric viruses.							
Targets	No. of gene copies tested	No. of replicates	No. of positive (%)				
Human-specific Bacteroides HF183 (HS-HF183) ^a	$10^3 - 10^2 - 10^1 - 10^0$	10	10 (100%) – 10 (100%) – 10 (100%) – 10 (100%)				
Human-specific enterococci surface protein (esp) marker (HS-esp)	$10^3 - 10^2 - 10^1 - 10^0$	10	10 (100%) – 10 (100%) – 10(100%) – 9 (90%)				
Human-specific adenoviruses (HS-AVs) ^a	$10^3 - 10^2 - 10^1 - 10^0$	10	10 (100%) – 10 (100%) – 10(100%) – 7 (70%)				
Human-specific polyomaviruses (HS-PVs) ^a	$10^3 - 10^2 - 10^1 - 10^0$	10	10 (100%) – 10 (100%) – 10(100%) – 3 (30%)				
Torquetenoviruses (TVs) ^a	$10^3 - 10^2 - 10^1 - 10^0$	10	10 (100%) – 10 (100%) – 10(100%) – 5 (50%)				
Enteroviruses (EVs) ^a	$10^3 - 10^2 - 10^1 - 10^0$	10	10 (100%) - 10 (100%) - 10(100%) - 1 (10%)				
Sapoviruses (SVs) ^a	$10^3 - 10^2 - 10^1 - 10^0$	10	10 (100%) - 10 (100%) - 10(100%) - 3 (30%)				
a Plasmid DNA.							

et al., 2008a; Carson et al., 2005; Gawler et al., 2007; Gourmelon et al., 2007). It has been reported that horizontal transfer of fecal bacteria may occur among species in close contact such as humans and their pets (Dick et al., 2005). A recent review recommended testing the specificity of the host-specific *Bacteroides* markers prior to application for MST field studies (Field and Samadpour, 2007).

The HS-esp markers were not detected in any animal fecal samples further supporting the high specificity of this marker

to distinguish between human and animal sources of fecal pollution. A recent study reported the presence of HS-*esp* markers in dog and gull fecal samples in the USA (Whitman et al., 2007). This could be explained by the fact that horizontal transfer of genes or cells is possible between humans and animals (Van den Boggard et al., 2002; Oancea et al., 2004). The HS-AVs and HS-PVs markers were not detected in any fecal or wastewater samples from the animals tested. The specificity of these markers has not been tested outside

Table 5 – Concentrations of fecal indicators and occurrence of host-specific molecular markers and enteric viruses in sewage spiked freshwater, seawater and distilled water samples.

Source water	Dilution	Concentrations of fecal indicators				No. of samples tested/real-time PCR positive results					
		E. coli	Enterococci	C. perfringens	HS-HF183	HS-esp	HS-AVs	HS-PVs	TVs	EVs	SVs
Freshwater	10^{-1}	$4.3\pm0.3\times10^{5}$	$9.6\pm0.2\times10^4$	$5.0\pm1.0\times10^2$	3/3	3/3	3/3	3/3	3/3	3/3	3/3
	10 ⁻²	$4.7\pm0.4\times10^4$	$8.3\pm0.6\times10^3$	$4.8\pm2.0\times10^{1}$	3/3	3/3	3/3	3/3	3/3	3/3	3/3
	10^{-3}	$4.1\pm0.6\times10^3$	$9.3\pm1.0\times10^2$	$5.1\pm3.0\times10^{0}$	3/3	3/3	3/3	3/3	3/3	3/3	3/3
	10^{-4}	$3.9\pm0.8\times10^2$	$8.1\pm6.0\times10^{1}$	ND	3/3	3/3	3/3	3/1	3/3	3/3	3/3
	10 ⁻⁵	$4.0\pm1.0\times10^{1}$	$7.1\pm3.0\times10^{0}$	ND	3/3	ND	ND	ND	ND	3/2	ND
	10 ⁻⁶	$3.0\pm2.0\times10^{0}$	ND	ND	3/3	ND	ND	ND	ND	ND	ND
	10^{-7}	ND	ND	ND	3/3	ND	ND	ND	ND	ND	ND
	10^{-8}	ND	ND	ND	3/2	ND	ND	ND	ND	ND	ND
	10^{-9}	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
	10^{-10}	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
Seawater	10^{-1}	$4.1\pm0.1\times10^{5}$	$9.8\pm0.4\times10^4$	$5.2\pm2.0\times10^2$	3/3	3/3	3/3	3/3	3/3	3/3	3/3
	10 ⁻²	$4.3\pm0.5\times10^4$	$9.1\pm0.3\times10^3$	$4.6\pm1.0\times10^{1}$	3/3	3/3	3/3	3/3	3/3	3/3	3/3
	10^{-3}	$4.0\pm0.8\times10^3$	$9.4\pm2.0\times10^2$	$8.1\pm6.0\times10^{0}$	3/3	3/3	3/3	3/2	3/3	3/3	3/3
	10^{-4}	$3.6\pm0.7\times10^2$	$7.1\pm9.0\times10^{1}$	ND	3/3	3/3	3/2	3/1	3/3	3/3	3/3
	10 ⁻⁵	$3.8\pm0.9\times10^{1}$	$6.1\pm6.0\times10^{0}$	ND	3/3	ND	ND	ND	ND	ND	ND
	10 ⁻⁶	$7.0\pm3.0\times10^{0}$	ND	ND	3/3	ND	ND	ND	ND	ND	ND
	10^{-7}	ND	ND	ND	3/3	ND	ND	ND	ND	ND	ND
	10^{-8}	ND	ND	ND	3/1	ND	ND	ND	ND	ND	ND
	10^{-9}	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
	10^{-10}	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
Distilled water	10^{-1}	$4.5\pm0.2\times10^{5}$	$9.9\pm0.2\times10^4$	$4.9\pm0.8\times10^2$	3/3	3/3	3/3	3/3	3/3	3/3	3/3
	10 ⁻²	$4.5\pm0.3\times10^4$	$8.8\pm0.6\times10^3$	$5.0\pm1.0\times10^{1}$	3/3	3/3	3/3	3/3	3/3	3/3	3/3
	10 ⁻²	$4.3\pm0.2\times10^3$	$9.1\pm3.0\times10^2$	$9.0\pm4.0\times10^{0}$	3/3	3/3	3/2	3/3	3/3	3/3	3/3
	10^{-4}	$4.1\pm0.5\times10^2$	$8.9\pm7.0\times10^{1}$	ND	3/3	3/3	3/1	3/2	3/3	3/3	3/3
	10 ⁻⁵	$3.6\pm0.7\times10^{1}$	$8.1\pm4.0\times10^{0}$	ND	3/3	ND	ND	ND	ND	ND	ND
	10^{-6}	$6.0\pm4.0\times10^{0}$	ND	ND	3/3	ND	ND	ND	ND	ND	ND
	10^{-7}	ND	ND	ND	3/3	ND	ND	ND	ND	ND	ND
	10^{-8}	ND	ND	ND	3/3	ND	ND	ND	ND	ND	ND
	10^{-9}	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
	10^{-10}	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

the USA and Spain where human-specific adenoviruses were shown to be highly sewage specific (Maluquer de Motes et al., 2004).

The HS-PVs can be shed in urine from a healthy individual and highly prevalent in sewage (McQuaig et al., 2006). However, limited data is available on the host-specificity of HS-PVs. In view of this, wastewater samples (i.e., mixture of urine and feces) were tested for host-specificity. To our knowledge, this is the first study that reports the host-specificity of HS-AVs and HS-PVs in sewage in Australia. All the markers tested in this study were present in all target samples except one sample which was negative for the HS-AVs marker. A limitation of the HS-esp, and viral markers is that their concentrations may be low in target samples and may not be present in fecal samples from all individuals (Scott et al., 2005; Field and Samadpour, 2007). Due to low concentrations, bacterial enrichment (for HS-esp), and nested realtime PCR protocols (for HS-AVs and HS-PVs) are required for the detection of these markers.

E. coli and enterococci were detected up to dilution 10^{-6} (contains 0.06 µl sewage) and 10^{-5} (contains 0.6 µl of sewage), respectively for freshwater, seawater, and distilled water samples spiked with sewage. However, HS-esp, HS-AVs, HS-PVs, and enteric viruses (i.e., SVs and TVs) were detected up to dilution 10^{-4} (contains 6.25 µl of sewage). EVs were detected up to dilution 10^{-5} due to the fact that this group of viruses consisted of polioviruses, coxsackieviruses and echoviruses, and therefore their concentrations are expected to be higher in sewage than other viruses tested. *C. perfringens* spores were detected up to dilution 10^{-3} (contains 62.5 µl of sewage). The low concentration of this indicator suggests that it may not be suitable to identify public health risks associated with sewage pollution in environmental waters.

The HS-HF183 markers were detected up to dilution 10^{-9} (contains 0.001 µl of sewage). However, at this dilution no culturable fecal indicators were detected. The number of HS-HF183F Bacteroides markers has been reported to be high, ranging from 2.0 \pm 0.2 \times 10 8 to 3.1 \pm 0.3 \times 10 9 per 100 ml raw sewage (Seurinck et al., 2005). Therefore, the real-time PCRs ability to detect these markers is expected to be high compared to the HS-esp and viral markers, as well as traditional fecal indicators. The HS-HF183 marker was detected in dilutions 3-4 orders of magnitude higher than those of enteric viruses implying that the concentration of this marker is much higher than enteric viruses in fresh sewage samples. From a public health point of view, such data needs to be interpreted with caution as a positive signal of the HS-HF183 marker in environmental waters may not always indicate the presence of enteric viruses or other pathogens. However, quantitative PCR could be used to quantify the HS-HF183 and enteric viruses in sewage samples to establish a relation between the concentration of the HS-HF183 and the occurrence of enteric viruses. For environmental samples, the HS-HF183 and HS-esp real-time PCR negative results do not rule out the presence of potential enteric pathogens and especially viruses which could have a different survival rate than these anaerobic and toxin gene markers.

It has to be noted that in this study, different DNA extraction methods and real-time PCR cycling parameters were used for different human-specific markers and enteric viruses. Therefore, it is possible that the real-time PCR may have underestimated the ability to detect fresh sewage for certain markers and/or enteric viruses. Nonetheless, the results at least indicate each published protocol's ability to detect fresh sewage in various matrices of water. The ability of the HS-HF183, HS-esp, HS-AVs, and HS-PVs markers to detect fresh sewage for different water matrices was similar. No discrepancies were observed probably due to the fact that fixed volume of fresh sewage was added into freshwater, seawater and distilled water matrices in the same manner and the spiked water samples were processed at the same time, and tested with the real-time PCR. The real-time PCR would detect target from both viable and non-viable cells. Therefore, the matrices did not have any effects on the realtime PCRs ability to detect host-specific markers and enteric viruses.

Limited data is available on the correlation between human-specific markers (i.e., Bacteroides and toxin gene) and enteric pathogens in environmental waters. A recent study reported the positive correlation between the HS-HF183 markers and bacterial zoonotic pathogens in environmental waters in the USA (Walters et al., 2007). Quantitative data on these markers would be required to obtain information regarding the likelihood of the presence of enteric viruses or other pathogens in environmental waters. Recently, quantitative real-time PCR methods have been developed to quantify human-specific Bacteroides and toxin gene markers in sewage and environmental waters (Ahmed et al., 2008c; Layton et al., 2006; Reischer et al., 2006; Seurinck et al., 2005). Consequently, these methods have the potential to provide information regarding the concentration of these markers, and the occurrence of pathogens in sewage and environmental waters.

The HS-esp, HS-AVs, and HS-PVs markers were detected up to dilution 10^{-4} , and the concentrations of E. coli and enterococci at this dilution were $3.9 \pm 0.8 \times 10^2$ and $8.1 \pm 6.0 \times 10^1$ CFUs respectively. It has to be noted that, at this dilution all the enteric viruses were also detected. The most important feature of the toxin gene and viral markers is that they provide direct information on pathogen status in environmental waters which could not be obtained using anaerobic markers or traditional fecal indicators. One potential drawback of toxin gene and viral markers is that their concentrations could be relatively low in environmental waters due to dilution and therefore, it may not be always possible to detect these markers. In this scenario, it is recommended that multiple markers (i.e., bacterial and viral) should be used in MST field studies to provide multiple lines of evidence of sewage pollution and public health risks.

5. Conclusions

• Human-specific bacterial and viral molecular markers tested in this study are highly sewage specific. To our knowledge this is the first study that evaluated the realtime PCR assays of four human-specific markers to detect fresh sewage in freshwater, seawater and distilled water matrices.

- The ability of the HS-HF183 marker to detect fresh sewage was 3–4 orders of magnitude higher than that of the HS-*esp* and viral markers. Based on our data, the HS-HF183 marker appears to be the most sensitive to detect fresh sewage pollution. However, its presence in environmental waters may not necessarily indicate the presence of enteric viruses due to their high abundance in sewage compared to enteric viruses.
- The ability of the real-time PCR assays of the humanspecific markers to detect fresh sewage in freshwater, seawater and distilled water matrices was similar suggesting that the matrices did not have any effects on the real-time PCR detection of fresh sewage.
- More research is required on the persistency of these markers in environmental water samples in relation to traditional fecal indicators and pathogenic microorganisms. In addition, quantitative real-time PCR data would be required to assess the magnitude of fecal pollution and associated public health risks. Our future research would focus on evaluating the survival of these markers in various environmental waters along with the traditional fecal indicators and pathogens.

Acknowledgements

This study was funded by Queensland Department of Natural Resources and Water (DNRW). This was a joint project between DNRW and Queensland University of Technology (QUT). We thank Dr. Flavia Huygens and Associate Professor Megan Hargreaves for providing laboratory space.

REFERENCES

- Ahmed, W., Tucker, J., Harper, J., Neller, R., Katouli, M., 2006. Comparison of the efficacy of an existing versus a locally developed metabolic fingerprint database to identify nonpoint sources of faecal contamination in a coastal lake. Water Res. 40 (12), 2339–2348.
- Ahmed, W., Stewart, J., Gardner, T., Powell, D., Brooks, P., Sullivan, D., Tindale, N., 2007. Sourcing faecal pollution: a combination of library-dependent and library-independent methods to identify human faecal pollution in non-sewered catchments. Water Res. 41 (16), 3771–3779.
- Ahmed, W., Stewart, J., Powell, D., Gardner, T., 2008a. Evaluation of Bacteroides markers for the detection of human faecal pollution. Lett. Appl. Microbiol. 46 (2), 237–242.
- Ahmed, W., Stewart, J., Powell, D., Gardner, T., 2008b. Evaluation of the host-specificity and prevalence of enterococci surface protein (*esp*) marker in sewage and its application for sourcing human faecal pollution. J. Environ. Qual. 37 (4), 1583–1588.
- Ahmed, W., Stewart, J., Powell, D., Gardner, T., 2008c. A real-time polymerase chain reaction assay for the quantitative detection of the human-specific enterococci surface protein marker in sewage and environmental waters. Environ. Microbiol. 10, 3255–3264.
- Bernhard, A.E., Field, K.G., 2000. A PCR assay to discriminate human and ruminant feces on the basis of host differences in Bacteroides–Prevotella genes encoding 16S rRNA. Appl. Environ. Microbiol. 66 (10), 4571–4574.

- Carson, C.A., Christiansen, J.M., Yampara-Iquise, H., Benson, V.W., Baffaut, C., Davis, J.V., Broz, R.R., Kurtz, W.B., Rogers, W.M., Fales, W.H., 2005. Specificity of a Bacteroides thetaiotaomicron marker for human feces. Appl. Environ. Microbiol. 71 (8), 4945–4949.
- Dick, L.K., Bernhard, A.E., Brodeur, T.J., Santo Domingo, J.W., Simpson, J.M., Walters, S.P., Field, K.G., 2005. Host distributions of uncultivated fecal *Bacteroidales* reveal genetic markers for fecal source identification. Appl. Environ. Microbiol. 71 (6), 3184–3191.
- Field, K.G., Samadpour, M., 2007. Fecal source tracking, the indicator paradigm, and managing water quality. Water Res. 41 (16), 3517–3538.
- Fong, T.T., Griffin, D.W., Lipp, E.K., 2005. Molecular assays for targeting human and bovine enteric viruses in coastal waters and their application for library-independent source tracking. Appl. Environ. Microbiol. 71 (4), 2070–2078.
- Fong, T.T., Lipp, E.K., 2005. Enteric viruses of humans and animals in aquatic environments: health risks, detection, and potential water quality assessment tools. Microbiol. Mol. Biol. Rev. 69 (2), 357–371.
- Gawler, A.H., Beecher, J.E., Brandão, J., Carroll, N.M., Falcão, L., Gourmelon, M., Masterson, B., Nunes, B., Porter, J., Rincé, A., Rodrigues, R., Thorp, M., Walters, J.M., Meijer, W.G., 2007. Validation of host-specific *Bacteriodales* 16S rRNA genes as markers to determine the origin of faecal pollution in Atlantic Rim countries of the European Union. Water Res. 41 (16), 3780–3784.
- Gourmelon, M., Caprais, M.P., Ségura, R., Le Mennec, C., Lozach, S., Piriou, J.Y., Rincé, A., 2007. Evaluation of two libraryindependent microbial source tracking methods to identify sources of fecal contamination in French estuaries. Appl. Environ. Microbiol. 73 (15), 4857–4866.
- Gordon, D.M., Bauer, S., Johnson, J.R., 2002. The genetic structure of Escherichia coli populations in primary and secondary habitats. Microbiology 148 (5), 1513–1522.
- Griffith, J.F., Weisburg, S.B., McGee, C.D., 2003. Evaluation of microbial source tracking methods using mixed fecal sources in aqueous test samples. J. Water Health 1 (4), 141–152.
- Hartel, P.G., Summer, J.D., Hill, J.L., Collins, J.V., Entry, J.A., Segars, W.I., 2002. Geographic variability of *Escherichia coli* ribotypes from animals in Idaho and Georgia. J. Environ. Qual. 31 (4), 1273–1278.
- Haramoto, E., Katayama, H., Oguma, K., Ohgaki, S., 2005. Application of cation-coated filter method to detection of noroviruses, enteroviruses, adenoviruses, and torqueteno viruses in the Tamagawa River in Japan. Appl. Environ. Microbiol. 71 (5), 2403–2411.
- Haramoto, E., Katayama, H., Phanwuan, C., Ohgaki, S., 2008. Quantitative detection of sapoviruses in wastewater and river in Japan. Lett. Appl. Microbiol. 46 (3), 408–413.
- Harwood, V.J., Wiggins, B.A., Hagedorn, C., Ellender, R.D.,
 Gooch, J., Kern, J., Samadpour, M., Chapman, A.C.H.,
 Robinson, B.J., 2003. Phenotypic library-based microbial source tracking methods: efficacy in the California collaborative study. J. Water Health 1 (4), 153–166.
- Hörman, A., Rimhanen-Finne, R., Maunula, L., von Bonsdorff, C.-H., Torvela, N., Heikinheimo, A., Hanninen, M.-L., 2004. *Campylobacter* spp., *Giardia* spp., *Cryptosporidium* spp., noroviruses, and indicator organisms in surface water in southwestern Finland, 2000–2001. Appl. Environ. Microbiol. 70 (1), 87–95.
- Khatib, L.A., Tsai, Y.L., Olson, B.H., 2002. A biomarker for the identification of cattle fecal pollution in water using the LTIIa toxin gene from enterotoxigenic *E. coli*. Appl. Microbiol. Biotechnol. 59 (1), 97–104.
- Layton, A., McKay, L., Williams, D., Garrett, V., Gentry, R., Sayler, G., 2006. Development of Bacteroides 16S rRNA gene TaqMan based real-time PCR assays for estimation of total,

human, and bovine fecal pollution in water. Appl. Environ. Microbiol. 72 (6), 4214–4224.

Maluquer de Motes, C., Clemente-Casares, P., Hundesa, A., Martin, M., Girones, R., 2004. Detection of bovine and porcine adenoviruses for tracing the source of fecal contamination. Appl. Environ. Microbiol. 70 (3), 1448–1454.

McQuaig, S.M., Scott, T.M., Harwood, V.J., Farrah, S.R., Lukasik, J.O., 2006. Detection of human-derived fecal pollution in environmental waters by use of a PCR-based human polyomavirus assay. Appl. Environ. Microbiol. 72 (12), 7567–7574.

Moore, D.F., Harwood, V.J., Ferguson, D.M., Lukasik, J., Hannah, P., Getrich, M., Brownell, M., 2005. Evaluation of antibiotic resistance analysis and ribotyping for identification of fecal pollution sources in an urban watershed. J. Appl. Microbiol. 99 (3), 618–628.

Oancea, C., Klare, I., Witte, W., Werner, G., 2004. Conjugative transfer of the virulence gene *esp*, among isolates of *Enterococcus faecium* and *Enterococcus faecalis*. J. Antimicrob. Chemother. 54 (1), 232–235.

Reischer, G.H., Kasper, D.C., Steinborn, R., Mach, R.L., Farnleitner, A.H., 2006. Quantitative PCR method for sensitive detection of ruminant fecal pollution in freshwater and evaluation of this method in Alpine Karstic regions. Appl. Environ. Microbiol. 72 (8), 5610–5614.

Santo Domingo, J.W., Bambic, D.G., Edge, T.A., Wuertz, S., 2007. Quo vadis source tracking? Towards a strategic framework for environmental monitoring of fecal pollution. Water Res. 41 (16), 3539–3552.

Scott, T.M., Rose, J.B., Jenkins, T.M., Farrah, S.R., Lukasik, J., 2002. Microbial source tracking: current methodology and future directions. Appl. Environ. Microbiol. 68 (12), 5796–5803.

Scott, T.M., Parveen, S., Portier, K.M., Rose, J.B., Tamplin, M.L., Farrah, S.A., Koo, A., Lukasik, J., 2003. Geographical variation in ribotype profiles of *Escherichia coli* isolates from humans, swine, poultry, beef, and dairy cattle in Florida. Appl. Environ. Microbiol. 69 (2), 1089–1092.

Scott, T.M., Jenkins, T.M., Lukasik, J., Rose, J.B., 2005. Potential use of a host-associated molecular in *Enterococcus faecium* as an index of human fecal pollution. Environ. Sci. Technol. 39 (1), 283–287.

- Seurinck, S., Defoirdt, T., Verstraete, W., Siciliano, S.D., 2005. Detection and quantification of the human-specific HF183 *Bacteroides* 16S rRNA genetic markers with real-time PCR for assessment of human fecal pollution in freshwater. Environ. Microbiol. 7 (2), 249–259.
- U.S. Environmental Protection Agency, 1997. Method 1600: Membrane Filter Test Method for Enterococci in Water. EPA/ 821/R-97/004. U.S. Environmental Protection Agency, Office of Water, Washington, DC.

U.S. Environmental Protection Agency, 2000. Improved Enumeration Methods for the Recreational Water Quality Indicators: Enterococci and Escherichia coli. Office of Science and Technology, Washington, DC, EPA/821/R-97/004, 55 pp.

U.S. Environmental Protection Agency, 2002. Method 1603: Escherichia coli (E. Coli) in Water by Membrane Filtration Using Modified Membrane-thermotolerant Escherichia coli Agar (Modified mTEC). EPA/821/R-02/023. U.S. Environmental Protection Agency, Office of Water, Washington, DC.

Van den Boggard, A.E., Willems, R., London, N., Top, J., Stobberingh, E.E., 2002. Antibiotic resistance of faecal enterococci in poultry, poultry farmers and poultry slayghterers. J. Antimicrob. Chemother. 49 (3), 497–505.

Walters, S.P., Gannon, V.P.G., Field, K.G., 2007. Detection of Bacteroidales fecal indicators and the zoonotic pathogens E. coli O157:H7, salmonella, and Campylobacter in river water. Environ. Sci. Technol. 41 (6), 1856–1862.

Whitman, R.L., Przzybyla-Kelly, K., Shivley, D.A., Byappanhali, M.N., 2007. Incidence of the enterococcal surface protein (*esp*) gene in human and animal faecal sources. Environ. Sci. Technol. 41 (17), 6090–6095.

Wiggins, B.A., 1996. Discriminant analysis of antibiotic resistance patterns in fecal streptococci, a method to differentiate human and animals sources of fecal pollution in natural waters. Appl. Environ. Microbiol. 62 (11), 3997–4002.

Wilson, I.G., 1997. Inhibition and facilitation of nucleic acid amplification. Appl. Environ. Microbiol. 63 (10), 3741–3751.

Wohlsen, T., Bayliss, J., Gray, B., Bates, J., Katouli, M., 2006. Evaluation of an alternative method for the enumeration and confirmation of Clostridium perfringens from treated and untreated sewage. Lett. Appl. Microbiol. 42 (5), 438–444.